Joshua M. Hall¹ and Daniel A. Warner¹

¹Department of Biological Sciences, Auburn University, Auburn, AL

jmh0131@auburn.edu

Embryo development and global change: how do reptile embryos respond to ecologically relevant thermal stress?


Abstract

Two components of global change, climate change and urbanization, both contribute to increased ambient temperatures that may induce heat stress or mortality in animals. Each phenomenon independently results in both increased mean temperatures and increased maximum day-time temperatures; however, there is also the potential for these components to act synergistically: extreme temperatures due to the urban heat island effect are likely to be exacerbated as the earth's surface warms due to climate change. Many animals can respond to harmful temperatures behaviorally, by altering their periods of activity or shifting their habitat use. Such behavioral compensation, however, is unavailable to embryos of ectotherms which typically develop inside eggs in the ground and receive little or no parental care. Thus, this early life stage is expected to be more vulnerable to harmful temperatures caused by aspects of global change, and yet, the effects of ecologically relevant thermal stress on these embryos has received little attention. We sought to understand the consequences of such extreme temperatures on embryological development by utilizing two species of lizard (*Anolis sagrei* and *Anolis cristatellus*) that commonly inhabit urban areas. We measured ground temperatures in an urban landscape where lizards nest and modeled daily thermal fluctuations that included brief periods of extremely high temperatures. We then subjected eggs of both species to various magnitudes and frequencies of these thermal fluctuations at multiple stages of embryo development. We report results of embryo survival and highlight the potential for extreme incubation temperatures to differentially impact species.



Embryo development and global change: how do reptile embryos respond to ecologically relevant thermal stress?

Joshua M Hall Daniel A Warner





Biodivers Conserv (2010) 19:1043–1062 DOI 10.1007/s10531-010-9782-4

ORIGINAL PAPER

Impacts of climate change on the amphibians and reptiles of Southeast Asia

David Bickford · Sam D. Howard · Daniel J. J. Ng · Jennifer A. Sheridan

Received: 31 August 2009/Accepted: 9 January 2010/Published online: 2 February 2010 © Springer Science+Business Media B.V. 2010

Biodivers Consery (2010) 19:1043-1062 DOI 10.1007/s10531-010-9782-4

ORIGINAL PAPER

Impacts of climate change on the amphibians and reptiles of Southeast Asia

David Bickford · Sam D. Howard · Daniel J. J. Ng · Jennifer A. Sheridan

Received: 31 August 2009/Accepted: 9 January 2010/Published online: 2 February 2010 © Springer Science+Business Media B.V. 2010

ECOLOGY

Are Lizards Toast?

Raymond B. Huey, 1 Jonathan B. Losos, 2 Craig Moritz3

evading thermal stress, tolerate high body temperatures, and resist water loss. Nevertheless, on page 894 of this issue, Sinervo et al. (1) document extinctions of lizobserved extinctions, to predict that warmpopulations extinct by 2080. If their prediction is even close to correct, lizards may be reproduction; extinction ensues. "the new amphibians" (2) in a race toward

A stark result for a genus of lizards in México leads off their paper: 12% of 200 previously validated Sceloporus populations (all with intact habitats) went extinct in recent

Department of Biology, University of Washington, Seattle, WA 98195, USA. 2 Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA. 3Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA. E-mail: hueyrb@uw.edu

izards should be relatively invulnera- decades. Moreover, extinction probability was ble to warming: They are very good at correlated with magnitude of warming at that site in spring, but not in other seasons. This correlation suggests that extinction is driven ard populations on five continents and argue rather than by summer heat stress. Lizard natthat global warming is responsible. They use ural history is instructive here: On hot days, a simple biological model, validated against lizards seek cooler refuges, such as burrows. With warming, lizards will spend longer periing will drive almost 40% of all global lizard ods in refuges, reducing foraging time, such

> To test this mechanistic hypothesis, Sinervo et al. examined four S. serrifer populamaximum available body temperatures of lizforcing retreat (see the second figure). At sites these data, the authors estimate that by 2080,

Warming is held responsible for a rash of extinctions of global lizard populations.

where the lizards are now extinct, predicted time restrictions exceeded 3.85 hours; but at sites where lizards persist, predicted restrictions were shorter. Sinervo et al. then used by energetic shortfalls during spring (when air temperature data from weather stations reproductive energy demands are highest), to estimate time restrictions at all Méxican sites. Sceloporus populations with predicted restrictions above 3.85 hours in spring had higher extinction rates than did populations with shorter restrictions.

To predict future extinctions, Sinervo et that net energy gain becomes insufficient for al. applied their history-validated approach to current and future warming scenarios across the globe, using 1216 lizard populations on four continents. First, by resurveying known tions, two of which have recently gone extinct lizard populations and conducting literature (see the first figure). Using field estimates of surveys, they detected many extinctions; for example, 21% of Madagascar lizard popuards (operative temperatures) at these sites lations in nature reserves have gone extinct. in spring and of body temperatures accept- Estimated activity-time restrictions (with critable for activity, they predicted the number ical thresholds tuned to the thermal biology of hours per day that operative temperatures of each lizard family) effectively predicted exceeded a lizard's thermal preferences, thus populations that had gone extinct. Based on

Biodivers Consery (2010) 19:1043-1062 DOI 10.1007/s10531-010-9782-4

ORIGINAL PAPER

Impacts of climate change on the amphibians and reptiles of Southeast Asia

David Bickford · Sam D. Howard · Daniel J. J. Ng · Jennifer A. Sheridan

Received: 31 Au © Springer Scie

Ecology Letters, (2011) 14: 484-492

doi: 10.1111/j.1461-0248.2011.01610.x

LETTER

Climate change threatens European conservation areas

/liquel B. Araújo, 1,2*↑ Diogo Jagador, 1,37 Mar Cabeza, 1,4 avid Nogués-Bravo^{1,5} and Vilfried Thuiller⁶

Europe has the world's most extensive network of conservation areas. Conservation areas are selected without taking into account the effects of climate change. How effectively would such areas conserve biodiversity under climate change? We assess the effectiveness of protected areas and the Natura 2000 network in conserving a large proportion of European plant and terrestrial vertebrate species under climate change. We found that by 2080, 58 ± 2.6% of the species would lose suitable climate in protected areas, whereas losses affected 63 ± 2.1% of the species of European concern occurring in Natura 2000 areas. Protected areas are expected to retain climatic suitability for species better than unprotected areas (P < 0.001), but Natura 2000 areas retain climate suitability for species no better and sometimes less effectively than unprotected areas. The risk is high that ongoing efforts to conserve Europe's biodiversity are jeopardized by climate change. New policies are required to avert this risk.

Keywords

Bioclimatic envelope models, climate change, conservation planning, gap analysis, Natura 2000 networks, protected areas.

Ecology Letters (2011) 14: 484-492

ECOLOGY

Are Lizards Toast?

Raymond B. Huev. Jonathan B. Losos. Craig Moritz

izards should be relatively invulnerable to warming: They are very good at evading thermal stress, tolerate high body temperatures, and resist water loss. Nevertheless, on page 894 of this issue, Sinervo et al. (1) document extinctions of lizard populations on five continents and argue that global warming is responsible. They use a simple biological model, validated against observed extinctions, to predict that warming will drive almost 40% of all global lizard populations extinct by 2080. If their prediction is even close to correct, lizards may be reproduction; extinction ensues. "the new amphibians" (2) in a race toward

A stark result for a genus of lizards in México leads off their paper: 12% of 200 previously validated Sceloporus populations (all with intact habitats) went extinct in recent

Department of Biology, University of Washington, Seattle, WA 98195, USA. 2 Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA. 3Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA. E-mail: hueyrb@uw.edu

decades. Moreover, extinction probability was correlated with magnitude of warming at that site in spring, but not in other seasons. This correlation suggests that extinction is driven by energetic shortfalls during spring (when reproductive energy demands are highest), rather than by summer heat stress. Lizard natural history is instructive here: On hot days, lizards seek cooler refuges, such as burrows. With warming, lizards will spend longer periods in refuges, reducing foraging time, such that net energy gain becomes insufficient for

To test this mechanistic hypothesis, Sinervo et al. examined four S. serrifer populations, two of which have recently gone extinct (see the first figure). Using field estimates of maximum available body temperatures of lizards (operative temperatures) at these sites in spring and of body temperatures acceptable for activity, they predicted the number exceeded a lizard's thermal preferences, thus forcing retreat (see the second figure). At sites these data, the authors estimate that by 2080,

Warming is held responsible for a rash of extinctions of global lizard populations.

where the lizards are now extinct, predicted time restrictions exceeded 3.85 hours; but at sites where lizards persist, predicted restrictions were shorter. Sinervo et al. then used air temperature data from weather stations to estimate time restrictions at all Méxican sites. Sceloporus populations with predicted restrictions above 3.85 hours in spring had higher extinction rates than did populations with shorter restrictions.

To predict future extinctions, Sinervo et al. applied their history-validated approach to current and future warming scenarios across the globe, using 1216 lizard populations on four continents. First, by resurveying known lizard populations and conducting literature surveys, they detected many extinctions; for example, 21% of Madagascar lizard populations in nature reserves have gone extinct. Estimated activity-time restrictions (with critical thresholds tuned to the thermal biology of hours per day that operative temperatures of each lizard family) effectively predicted populations that had gone extinct. Based on

Biodivers Consery (2010) 19:1043-1062 DOI 10.1007/s10531-010-9782-4

ORIGINAL PAPER

Impacts of climate change on the amphibians and reptiles of Southeast Asia

David Bickford · Sam D. Howard · Daniel J. J. Ng · Jennifer A. Sheridan

Received: 31 Au © Springer Scie

Ecology Letters, (2011) 14: 484-492

doi: 10.1111/j.1461-0248.2011.01610.x

LETTER

Climate change threatens European conservation areas

/liquel B. Araújo, 1,2*↑ Diogo Jagador, 1,37 Mar Cabeza, 1,4 avid Nogués-Bravo^{1,5} and Vilfried Thuiller⁶

Europe has the world's most extensive network of conservation areas. Conservation areas are selected without taking into account the effects of climate change. How effectively would such areas conserve biodiversity under climate change? We assess the effectiveness of protected areas and the Natura 2000 network in conserving a large proportion of European plant and terrestrial vertebrate species under climate change. We found that by 2080, 58 ± 2.6% of the species would lose suitable climate in protected areas, whereas losses affected 63 ± 2.1% of the species of European concern occurring in Natura 2000 areas. Protected areas are expected to retain climatic suitability for species better than unprotected areas (P < 0.001), but Natura 2000 areas retain climate suitability for species no better and sometimes less effectively than unprotected areas. The risk is high that ongoing efforts to conserve Europe's biodiversity are jeopardized by climate change. New policies are required to avert this risk.

Keywords

Bioclimatic envelope models, climate change, conservation planning, gap analysis, Natura 2000 networks, protected areas.

Ecology Letters (2011) 14: 484-492

Erosion of Lizard Diversity by Climate Change and Altered Thermal Niches

Barry Sinervo. 1,15* Fausto Méndez-de-la-Cruz. Donald B. Miles. 3,15 Benoit Heulin. 4 Elizabeth Bastiaans, 1 Maricela Villagrán-Santa Cruz, 5 Rafael Lara-Resendiz, 2 Norberto Martínez-Méndez, Martha Lucía Calderón-Espinosa, Rubi Nelsi Meza-Lázaro, 2 Héctor Gadsden, Luciano Javier Avila, Mariana Morando, Ignacio J. De la Riva, 9 Pedro Victoriano Sepulveda, 10 Carlos Frederico Duarte Rocha, 11 Nora Ibargüengoytía, 12 César Aquilar Puntriano. 13 Manuel Massot. 14 Virginie Lepetz. 15 † Tuula A. Oksanen. 16 David G. Chapple, 17 Aaron M. Bauer, 18 William R. Branch, 19 Jean Clobert, 15 Jack W. Sites Jr. 20

It is predicted that climate change will cause species extinctions and distributional shifts in coming decades, but data to validate these predictions are relatively scarce. Here, we compare recent and historical surveys for 48 Mexican lizard species at 200 sites. Since 1975, 12% of local populations have gone extinct. We verified physiological models of extinction risk with observed local extinctions and extended projections worldwide. Since 1975, we estimate that 4% of local populations have gone extinct worldwide, but by 2080 local extinctions are projected to reach 39% worldwide, and species extinctions may reach 20%. Global extinction projections were validated with local extinctions observed from 1975 to 2009 for regional biotas on four other continents, suggesting that lizards have already crossed a threshold for extinctions caused by climate change.

I label alimete abanca offects arganisms (I) but not autinations (II) Hance there is still

ECOLOGY

Are Lizards Toast?

Raymond B. Huev. Jonathan B. Losos. 2 Craig Moritz3

izards should be relatively invulnerable to warming: They are very good at ✓ evading thermal stress, tolerate high body temperatures, and resist water loss. Nevertheless, on page 894 of this issue, Sinervo et al. (1) document extinctions of lizard populations on five continents and argue that global warming is responsible. They use a simple biological model, validated against observed extinctions, to predict that warming will drive almost 40% of all global lizard populations extinct by 2080. If their prediction is even close to correct, lizards may be reproduction; extinction ensues. "the new amphibians" (2) in a race toward

A stark result for a genus of lizards in México leads off their paper: 12% of 200 previously validated Sceloporus populations (all with intact habitats) went extinct in recent

Department of Biology, University of Washington, Seattle, WA 98195, USA. 2 Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA, 3Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA. E-mail: hueyrb@uw.edu

decades. Moreover, extinction probability was correlated with magnitude of warming at that site in spring, but not in other seasons. This correlation suggests that extinction is driven by energetic shortfalls during spring (when reproductive energy demands are highest), rather than by summer heat stress. Lizard natural history is instructive here: On hot days, lizards seek cooler refuges, such as burrows. With warming, lizards will spend longer periods in refuges, reducing foraging time, such that net energy gain becomes insufficient for

To test this mechanistic hypothesis, Sinervo et al. examined four S. serrifer populations, two of which have recently gone extinct (see the first figure). Using field estimates of maximum available body temperatures of lizards (operative temperatures) at these sites in spring and of body temperatures acceptable for activity, they predicted the number of hours per day that operative temperatures exceeded a lizard's thermal preferences, thus forcing retreat (see the second figure). At sites these data, the authors estimate that by 2080,

Warming is held responsible for a rash of extinctions of global lizard populations.

where the lizards are now extinct, predicted time restrictions exceeded 3.85 hours; but at sites where lizards persist, predicted restrictions were shorter. Sinervo et al. then used air temperature data from weather stations to estimate time restrictions at all Méxican sites. Sceloporus populations with predicted restrictions above 3.85 hours in spring had higher extinction rates than did populations with shorter restrictions.

To predict future extinctions, Sinervo et al. applied their history-validated approach to current and future warming scenarios across the globe, using 1216 lizard populations on four continents. First, by resurveying known lizard populations and conducting literature surveys, they detected many extinctions; for example, 21% of Madagascar lizard populations in nature reserves have gone extinct. Estimated activity-time restrictions (with critical thresholds tuned to the thermal biology of each lizard family) effectively predicted populations that had gone extinct. Based on

Biodivers Consery (2010) 19:1043-1062 DOI 10.1007/s10531-010-9782-4

ORIGINAL PAPER

Impacts of climate change on the amphibians and rept Wilfried Thuiller of Southeast Asia

David Bickford · Sam D. Howard · Daniel J. J. Ng · Jennifer A. Sherida



Climate change and the ecologist

The evidence for rapid climate change now seems overwhelming. Global temperatures are predicted to rise by up to 4 °C by 2100, with associated alterations in precipitation patterns. Assessing the consequences for biodiversity, and how they might be mitigated, is a Grand Challenge in ecology.

How serious is climate change compared with other factors affecting biodiversity?

Very - but it tends to act over a longer timescale. The ecological disruption wrought

the Northern Hemisphere may have to move northward by some 500 km (or 500 m higher in altitude) to find a suitable climatic regime. Higher temperatures are likely to be accompanied by more humid, wetter conditions, but

photosynthetic pathway or 'woodiness', as well as the subterranean organisms associated with them. Likewise, an increase of anthropogenic atmospheric nitrogen deposition affects nitrogen-limited regions (temperate and boreal

Erosion of Lizard Diversity by Climate Change and Altered Thermal Niches

Barry Sinervo. 1,15* Fausto Méndez-de-la-Cruz. Donald B. Miles. 3,15 Benoit Heulin. 4 Elizabeth Bastiaans, 1 Maricela Villagrán-Santa Cruz, 5 Rafael Lara-Resendiz, 2 Norberto Martínez-Méndez, Martha Lucía Calderón-Espinosa, Rubi Nelsi Meza-Lázaro, 2 or Gadsden, Luciano Javier Avila, Mariana Morando, Ignacio J. De la Riva, 9 o Victoriano Sepulveda, 10 Carlos Frederico Duarte Rocha, 11 Nora Ibargüengoytía, 12 r Aguilar Puntriano. 13 Manuel Massot. 14 Virginie Lepetz. 15† Tuula A. Oksanen. 16 d G. Chapple, ¹⁷ Aaron M. Bauer, ¹⁸ William R. Branch, ¹⁹ Jean Clobert, ¹⁵ Jack W. Sites Jr. ²⁰

predicted that climate change will cause species extinctions and distributional shifts in coming des, but data to validate these predictions are relatively scarce. Here, we compare recent historical surveys for 48 Mexican lizard species at 200 sites. Since 1975, 12% of local lations have gone extinct. We verified physiological models of extinction risk with observed local ctions and extended projections worldwide, Since 1975, we estimate that 4% of local lations have gone extinct worldwide, but by 2080 local extinctions are projected to reach 39% lwide, and species extinctions may reach 20%. Global extinction projections were validated local extinctions observed from 1975 to 2009 for regional biotas on four other continents, esting that lizards have already crossed a threshold for extinctions caused by climate change.

I lobal climate change effects arganisms (I) but not outlingtions (I). House there is still

Received: 31 Au © Springer Scie

Ecology Letters, (2011) 14: 484-492

doi: 10.1111/j.1461-0248.2011.01610.x

LETTER

Climate change threatens European conservation areas

/liquel B. Araújo, 1,2*↑ Diogo Jagador, 1,37 Mar Cabeza, 1,4 avid Nogués-Bravo^{1,5} and Vilfried Thuiller⁶

Europe has the world's most extensive network of conservation areas. Conservation areas are selected without taking into account the effects of climate change. How effectively would such areas conserve biodiversity under climate change? We assess the effectiveness of protected areas and the Natura 2000 network in conserving a large proportion of European plant and terrestrial vertebrate species under climate change. We found that by 2080, 58 ± 2.6% of the species would lose suitable climate in protected areas, whereas losses affected 63 ± 2.1% of the species of European concern occurring in Natura 2000 areas. Protected areas are expected to retain climatic suitability for species better than unprotected areas (P < 0.001), but Natura 2000 areas retain climate suitability for species no better and sometimes less effectively than unprotected areas. The risk is high that ongoing efforts to conserve Europe's biodiversity are jeopardized by climate change. New policies are required to avert this risk.

Keywords

Bioclimatic envelope models, climate change, conservation planning, gap analysis, Natura 2000 networks, protected areas.

Ecology Letters (2011) 14: 484-492

ECOLOGY

Are Lizards Toast?

Raymond B. Huev. Jonathan B. Losos. Craig Moritz

izards should be relatively invulnerable to warming: They are very good at ✓ evading thermal stress, tolerate high body temperatures, and resist water loss. Nevertheless, on page 894 of this issue, Sinervo et al. (1) document extinctions of lizard populations on five continents and argue that global warming is responsible. They use a simple biological model, validated against observed extinctions, to predict that warming will drive almost 40% of all global lizard populations extinct by 2080. If their prediction is even close to correct, lizards may be "the new amphibians" (2) in a race toward

A stark result for a genus of lizards in México leads off their paper: 12% of 200 previously validated Sceloporus populations (all with intact habitats) went extinct in recent

Department of Biology, University of Washington, Seattle, WA 98195, USA. 2 Museum of Comparative Zoology, Harvard University, Cambridge, MA 02138, USA, 3Museum of Vertebrate Zoology, University of California, Berkeley, CA 94720, USA. E-mail: hueyrb@uw.edu

decades. Moreover, extinction probability was correlated with magnitude of warming at that site in spring, but not in other seasons. This correlation suggests that extinction is driven by energetic shortfalls during spring (when reproductive energy demands are highest), rather than by summer heat stress. Lizard natural history is instructive here: On hot days, lizards seek cooler refuges, such as burrows. With warming, lizards will spend longer periods in refuges, reducing foraging time, such that net energy gain becomes insufficient for reproduction; extinction ensues.

To test this mechanistic hypothesis, Sinervo et al. examined four S. serrifer populations, two of which have recently gone extinct (see the first figure). Using field estimates of maximum available body temperatures of lizards (operative temperatures) at these sites in spring and of body temperatures acceptable for activity, they predicted the number of hours per day that operative temperatures exceeded a lizard's thermal preferences, thus forcing retreat (see the second figure). At sites


Warming is held responsible for a rash of extinctions of global lizard populations.


where the lizards are now extinct, predicted time restrictions exceeded 3.85 hours; but at sites where lizards persist, predicted restrictions were shorter. Sinervo et al. then used air temperature data from weather stations to estimate time restrictions at all Méxican sites. Sceloporus populations with predicted restrictions above 3.85 hours in spring had higher extinction rates than did populations with shorter restrictions.


To predict future extinctions, Sinervo et al. applied their history-validated approach to current and future warming scenarios across the globe, using 1216 lizard populations on four continents. First, by resurveying known lizard populations and conducting literature surveys, they detected many extinctions; for example, 21% of Madagascar lizard populations in nature reserves have gone extinct. Estimated activity-time restrictions (with critical thresholds tuned to the thermal biology of each lizard family) effectively predicted populations that had gone extinct. Based on these data, the authors estimate that by 2080,


Global Change - urbanization
















0 0.5 1.0 1.5 2.0 2.5 3.0

 Average temperatures
 higher in urban areas than natural sites




0 0.5 1.0 1.5 2.0 2.5 3.0

- Average temperatures
 higher in urban areas than natural sites
- Thermal fluctuations,
 thermal spikes, more
 extreme in urban areas




0 0.5 1.0 1.5 2.0 2.5 3.0


- Average temperatures
 higher in urban areas than natural sites
- Thermal fluctuations,
 thermal spikes, more
 extreme in urban areas
- Climate change in 'real time'



- Average temperatures
 higher in urban areas than natural sites
- Thermal fluctuations,
 thermal spikes, more
 extreme in urban areas
- Climate change in 'real time'
- Implications for ectotherms (e.g. reptiles)



- Average temperatures
 higher in urban areas than natural sites
- Thermal fluctuations,
 thermal spikes, more
 extreme in urban areas
- Climate change in 'real time'
- Implications for ectotherms (e.g. reptiles)
 - Adults




- Average temperatures
 higher in urban areas than natural sites
- Thermal fluctuations,
 thermal spikes, more
 extreme in urban areas
- Climate change in 'real time'
- Implications for ectotherms (e.g. reptiles)
 - Adults
 - Developing embryos?



 Are particularly sensitive to environmental disturbance (no parental care)



- Are particularly sensitive to environmental disturbance (no parental care)
- Are unable to behaviorally compensate for adverse conditions (can't run away)



- Are particularly sensitive to environmental disturbance (no parental care)
- Are unable to behaviorally compensate for adverse conditions (can't run away)
- Influence population dynamics and species distributions and persistence




Pike. 2014. Glob Chang Biol 20: 7–15

• Not Ecologically relevant

- Not Ecologically relevant
 - Heat shocks


- Not Ecologically relevant
 - Heat shocks
 - Constant temperature incubation

- Not Ecologically relevant
 - Heat shocks
 - Constant temperature incubation



- Not Ecologically relevant
 - Heat shocks
 - Constant temperature incubation







- Not Ecologically relevant
 - Heat shocks
 - Constant temperature incubation




Reduce egg survival by ~ 20%

Hall & Warner. 2017. Biol J Linn Soc Hall & Warner. 2018. J Exp Biol







Magnitude (how hot?)




- Magnitude (how hot?)
- Frequency (how often?)



- Magnitude (how hot?)
- Frequency (how often?)
- Timing (how old?)




- Magnitude (how hot?)
- Frequency (how often?)
- Timing (how old?)



Ecologically relevant thermal stress

- Magnitude (how hot?)
- Frequency (how often?)
- Timing (how old?)
- Species (how general?)
























































































Embryo survival




Embryo survival




Treatment: χ^2_1 =58.46; p<0.0001

Treatment by Temperature: χ^2_1 =51.52; p<0.0001




Peak temperature (°C)

Embryo survival




Treatment: χ^2_1 =58.46; p<0.0001

Treatment by Temperature: χ^2_1 =51.52; p<0.0001




Treatment: χ^2_1 =20.45; p<0.0001

Treatment by Temperature: χ^2_1 =62.7; p<0.0001




Treatment: χ^2_1 =58.46; p<0.0001


Treatment by Temperature: χ^2_1 =51.52; p<0.0001





Treatment: χ^2_1 =20.45; p<0.0001

Treatment by Temperature: χ^2_1 =62.7; p<0.0001











Age: χ^2_1 =5.51; p=0.02


Age by Treatment: χ^2_1 =5.14; p=0.02





Age: χ^2_1 =5.51; p=0.02


Age by Treatment: χ^{2}_{1} =5.14; p=0.02




Age: χ^2_1 =3.63; p=0.06


Age by Treatment: χ^2_1 =4.87; p=0.03

Why are older embryos less robust?



Why are older embryos less robust?






Thompson & Stewart; CBP 1997


Why are older embryos less robust?







Thompson & Stewart; CBP 1997

 Thermal tolerance of embryos may vary widely across species







- Thermal tolerance of embryos may vary widely across species
- Tolerance changes through development



Ollonen et al. 2018. Front. Physiol.





- Thermal tolerance of embryos may vary widely across species
- Tolerance changes through development


- Thermal tolerance of embryos may vary widely across species
- Tolerance changes through development



Thermal tolerances of sea turtle embryos: current understanding and future directions

Robert Howard¹, Ian Bell², David A. Pike^{1,*}

¹School of Marine and Tropical Biology, James Cook University, Townsville, Queensland 4811, Australia ²Department of Environment and Heritage Protection, Townsville, Queensland 4814, Australia



- Thermal tolerance of embryos may vary widely across species
- Tolerance changes through development



Thermal tolerances of sea turtle embryos: current understanding and future directions

Robert Howard¹, Ian Bell², David A. Pike^{1,*}

¹School of Marine and Tropical Biology, James Cook University, Townsville, Queensland 4811, Australia
²Department of Environment and Heritage Protection, Townsville, Queensland 4814, Australia

ture range that sea turtle embryos can withstand and at which they can successfully hatch, but have not yet determined whether temperature fluctuation and stage of embryonic development interact, such that the thermal tolerance of embryos changes during incubation. Research on olive ridley sea turtles *Lepido*-

Acknowledgements

Mentors


Dan Warner Tim Mitchell Renata Brandt

<u>Undergraduates</u>

Austin Hulbert Kayla Wilson Ryan Weesner David Quinn Chanel Reali

Collaborators

Jason Kolbe James Stroud





Funding

National Science Foundation

Acknowledgements

Mentors

Dan Warner Tim Mitchell Renata Brandt

<u>Undergraduates</u>

Austin Hulbert Kayla Wilson Ryan Weesner David Quinn Chanel Reali

Collaborators

Jason Kolbe James Stroud

Funding

National Science Foundation







Literature Cited

Hall, J.M. and Warner, D.A., 2017. Body size and reproduction of a non-native lizard are enhanced in an urban environment. *Biological Journal of the Linnean Society*, 122(4), pp.860-871.

Hall, J.M. and Warner, D.A., 2018. Thermal spikes from the urban heat island increase mortality and alter physiology of lizard embryos. *Journal of Experimental Biology*, 221(14), p.jeb181552.

Howard, R., Bell, I. and Pike, D.A., 2014. Thermal tolerances of sea turtle embryos: current understanding and future directions. *Endangered Species Research*, 26(1), pp.75-86.

Pike, D.A., 2014. Forecasting the viability of sea turtle eggs in a warming world. *Global change biology*, 20(1), pp.7-15.