

*Performance
of Soybean
Varieties in
Alabama,
2012*

*Agronomy and Soils Departmental Series No. 327
Alabama Agricultural Experiment Station
William Batchelor, Director
Auburn University, Auburn, Alabama,
December 2012*

*Printed in cooperation with the Alabama Cooperative Extension System
(Alabama A&M University and Auburn University)*

| | |
|---|----|
| Table 18. Performance of Group IV and V Soybean Varieties on Sumter Soil, Marion Junction, Alabama, 2012 | 23 |
| Table 19. Performance of Group IV and V Soybean Varieties on Sumter Soil, Marion Junction, Alabama. Three-year Summary, 2010 - 2012..... | 24 |
| Table 20. Performance of Group VI and VII Soybean Varieties on Vaiden Soil, Marion Junction, Alabama, 2012 | 25 |
| Table 21. Performance of Group VI and VII Soybean Varieties on Vaiden Soil, Marion Junction, Alabama, 2012, Three-year Summary, 2010 - 2012..... | 26 |
| Table 22. Performance of Group VI and VII Soybean Varieties on Sumter Soil, Marion Junction, Alabama, 2012 | 27 |
| Table 23. Performance of Group VI and VII Soybean Varieties on Sumter Soil, Marion Junction, Alabama. Three-year Summary, 2010 - 2012..... | 28 |
| SOUTHERN ALABAMA | |
| Table 24. Performance of Group IV and V Soybean Varieties at Fairhope, Alabama, 2012 | 29 |
| Table 25. Performance of Group IV and V Soybean Varieties at Fairhope, Alabama, Three-year Summary, 2010 - 2012 | 30 |
| Table 26. Performance of Group VI and VII Soybean Varieties at Fairhope, Alabama, 2012 | 31 |
| Table 27. Performance of Group VI and VII Soybean Varieties at Fairhope, Alabama, Three-year Summary, 2010 - 2012 | 32 |
| Table 28. Performance of Group IV and V Soybean Varieties at Brewton, Alabama, 2012..... | 33 |
| Table 29. Performance of Group IV and V Soybean Varieties at Brewton, Alabama, Three-year Summary, 2010 and 2012 | 34 |
| Table 30. Performance of Group VI and VII Soybean Varieties at Brewton, Alabama, 2012..... | 35 |
| Table 31. Performance of Group VI and VII Soybean Varieties at Brewton, Alabama, Three-year Summary, 2010 and 2012 | 36 |
| ANCILLARY INFORMATION | |
| Table 32. Cultural Practices for Soybean Variety Tests in 2012..... | 37 |
| Table 33. Soil Types for Soybean Tests, 2012 | 38 |
| Table 34. Rainfall at Test Locations During Growing Season, 2012 | 39 |
| Table 35. Entries and Sources of Seed for Soybean Tests, 2012..... | 40 |

PERFORMANCE OF SOYBEAN VARIETIES IN ALABAMA, 2012

K. M. Glass, D.P. Delaney, and Edzard van Santen

Advisor, Natl. Res. Prog., Extension Soybean Specialist, and Professor

INTRODUCTION

Soybean variety tests are conducted annually by the Alabama Agricultural Experiment Station. The 7 locations used represent the major soil and climatic regions of Alabama. These locations are divided into logical soybean growing regions. The regions and locations are:

| Region | Location |
|------------|---------------------------|
| Northern | Belle Mina, Crossville |
| Central | Tallassee, Shorter |
| Southern | Brewton |
| Black Belt | Marion Junction (2 soils) |
| Gulf Coast | Fairhope |

EXPERIMENTAL PROCEDURES

The standard tests were conducted as a randomized complete block design with four replications. Standard plot size was four 30- to 38-inch rows by 20 feet long. Fifteen feet of the middle two rows were harvested for yield. Seeding rate was 10 viable seeds per foot of row. The Group IV test was drilled with seven 7-inch rows. Seeding rate was five viable seeds per foot of row.

Data were collected on seed yield, moisture, lodging, shattering, plant height, and maturity date. Plot yields were adjusted to 13 percent moisture and converted to bushels (60 pounds) per acre. Lodging was scored on a scale of 1 to 5 as follows:

- 1 - almost all plants erect.
- 2 - either all plants leaning slightly (less than 45%) or a few plants down.
- 3 - either all plants leaning moderately (approximately 45%) or 25 to 50 percent of the plants down.
- 4 - either all plants leaning more than 45% or 50 to 80 percent of the plants down.
- 5 - more than 80 percent of the plants down.

Shattering was rated on a scale of 1 to 5 based on performance of the border rows 14 days after maturity. A rating of 1 indicates no shattering, a rating of 3 indicates a 4 to 8 percent shattering, and a rating of 5 is 20 percent or more shattering. Plant height was determined by measuring from the ground to the top of the plant at maturity. Maturity date was the day 95 percent of the pods achieved mature pod color. Harvest was approximately 7 to 10 days later.

SEASONAL CONDITIONS

Rainfall for 2012 is shown in Table 30. The normal planting dates for the standard tests are the first week in May, May 15-25, and May 25 to June 5 for northern, central, and southern Alabama locations, respectively.

COMPARING VARIETIES

To aid in determining real yield differences, a statistical analysis of variance was performed on the data from each location. The L.S.D. (least significant difference) and C.V. (coefficient of variation) are reported for each location's 2012 test, and for the location's or region's 2- and 3-year averages. The difference in yield of two varieties must exceed the L.S.D. value for one variety to be considered superior to others in yield in that particular test. The C.V. is a measure of the variability in an experiment. An increase in its value indicates an increase in the unexplained variability.

Since the performance of varieties varies with location and year, long-term averages from several locations are more reliable than 1-year performance. Three-year regional averages are considered a reliable evaluation of the relative performance of varieties. However varietal rankings may change among years and among locations. This change in rankings is measured by the significance of variety x location, variety x year, variety x location*year interaction. These interactions were significant in all cases. Thus, care should be exercised when extrapolating results from one location or year to another.

ACKNOWLEDGMENTS

Appreciation is expressed to the following station superintendents and their staffs. It is their quality work, which makes this report a reliable source of information for farmers in their regions.

Chet Norris and David Harkins, Tennessee Valley Research and Extension Center; Joyce Ducar, Sand Mountain Research and Extension Center; Steve Nightengale, E.V. Smith Research Center, Plant Breeding Unit; Shawn Scott, E.V. Smith Research Center, Field Crops Research Unit; Jimmy Holliman, Black Belt Research and Extension Center; Randy Akridge, Brewton Agricultural Research Unit; Malcomb Pegues and Jarrod Jones, Gulf Coast Research and Extension Center.

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 1. PERFORMANCE OF GROUP IV SOYBEAN VARIETIES IN NORTHERN ALABAMA, 2012

| Variety | Belle Mina | Cross ville | Regional Average | | | | |
|---------------------|---------------------|----------------|------------------|------------------|---------------------|-----------------|-------------|
| | | | Yield | Lodging score | Shattering score | Plant height | |
| | ----- bu/acre ----- | | - inches - | | | | |
| Maturity Group IV | | | | | | | |
| Dyna-Gro S48RS53 | 64.0 | 75.4 | 69.7 | 1.7 | 1.6 | 34 | 9-17 |
| HBK RY4620 | 62.1 | 76.4 | 69.2 | 1.8 | 1.5 | 29 | 9-20 |
| USG 74A92R | 72.6 | 63.2 | 67.9 | 2.0 | 1.3 | 33 | 9-26 |
| HBK RY4721 | 62.9 | 70.1 | 66.5 | 1.5 | 1.5 | 35 | 9-17 |
| Schillinger 4990.RC | 66.0 | 65.1 | 65.6 | 1.5 | 1.4 | 34 | 9-24 |
| Asgrow AG 4632 | 61.3 | 69.6 | 65.5 | 1.5 | 1.3 | 32 | 9-18 |
| Asgrow AG 4832 | 65.4 | 64.4 | 64.9 | 2.1 | 1.8 | 35 | 9-19 |
| Dyna-Gro 37RY47 | 59.0 | 70.2 | 64.6 | 1.1 | 1.5 | 28 | 9-15 |
| USG 74G99L | 69.5 | 59.2 | 64.3 | 1.4 | 1.1 | 35 | 9-25 |
| USG 74E88 | 64.4 | 63.2 | 63.8 | 1.4 | 1.7 | 33 | 9-13 |
| Terral REV 49R43 | 58.0 | 66.6 | 62.3 | 1.6 | 1.3 | 30 | 9-21 |
| GoSoy 4910LL | 62.3 | 61.6 | 62.0 | 1.4 | 1.3 | 33 | 9-29 |
| HBK R4924 | 66.1 | 57.3 | 61.7 | 1.7 | 1.1 | 37 | 9-19 |
| Schillinger 495.RC | 67.3 | 56.1 | 61.7 | 2.0 | 2.1 | 35 | 9-20 |
| USG 74H92R | 65.6 | 56.5 | 61.0 | 1.5 | 1.0 | 29 | 9-21 |
| Dyna-Gro S47RY13 | 62.3 | 57.2 | 59.7 | 1.3 | 1.8 | 30 | 9-17 |
| Terral REV 47R53 | 55.6 | 63.2 | 59.4 | 1.8 | 1.1 | 31 | 9-15 |
| TN09-029 | 61.7 | 56.8 | 59.3 | 1.1 | 1.0 | 26 | 9-24 |
| Terral REV 48R33 | 55.2 | 63.1 | 59.2 | 1.8 | 1.5 | 33 | 9-12 |
| GoSoy 4411LL | 58.8 | 58.9 | 58.9 | 1.8 | 3.0 | 32 | 9-15 |
| Terral REV 49R22 | 63.5 | 49.6 | 56.5 | 1.9 | 2.6 | 32 | 9-21 |
| Asgrow AG 4633 | 61.1 | 47.2 | 54.1 | 1.3 | 1.6 | 26 | 9-12 |
| Terral REV 47R74 | 55.5 | 52.7 | 54.1 | 1.1 | 2.6 | 28 | 9-14 |
| Asgrow AG 4933 | 64.3 | 43.7 | 54.0 | 1.1 | 1.3 | 30 | 9-20 |
| Terral REV 49R54 | 55.3 | | | | | | 9-19 |
| Trial mean | 62.4 | 61.1 | 61.9 | 1.6 | 1.6 | 32 | 9-19 |
| LSD(0.10) | 5.0 | 7.8 | 4.8 | | | | |
| CV (%) | 7.8 | 12.6 | 10.8 | | | | |

PERFORMANCE OF SOYBEAN VARIETIES IN ALABAMA, 2012

7

TABLE 2. PERFORMANCE OF GROUP IV SOYBEAN VARIETIES IN NORTH ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | | | | Maturity date |
|---------------------|---------------------|-----------------------|-----------------------|---------------|------------------|--------------|---------------|
| | 2012 | 2-yr avg [†] | 3-yr avg [†] | Lodging score | Shattering score | Plant height | |
| | ----- bu/acre ----- | | | | | | - inch - |
| Maturity Group IV | | | | | | | |
| Schillinger 495.RC | 61.7 | 56.9 | 46.9 | 1.7 | 2.2 | 36 | 9-10 |
| Schillinger 4990.RC | 65.6 | 59.0 | 46.9 | 1.3 | 1.9 | 34 | 9-12 |
| Terral REV 49R22 | 56.5 | 54.0 | 45.5 | 1.7 | 2.3 | 34 | 9-8 |
| USG 74E88 | 63.8 | 63.9 | | 1.6 | 1.8 | 35 | 9-9 |
| Dyna-Gro 37RY47 | 64.6 | 63.7 | | 1.2 | 1.6 | 29 | 9-14 |
| Terral REV 48R33 | 59.2 | 63.4 | | 2.2 | 1.6 | 35 | 9-9 |
| Terral REV 49R43 | 62.3 | 62.7 | | 1.7 | 1.3 | 31 | 9-16 |
| Terral REV 47R53 | 59.4 | 59.3 | | 1.7 | 1.3 | 32 | 9-12 |
| Dyna-Gro S48RS53 | 69.7 | | | 1.7 | 1.6 | 34 | 9-17 |
| HBK RY4620 | 69.2 | | | 1.9 | 1.6 | 29 | 9-20 |
| USG 74A92R | 67.9 | | | 2.0 | 1.3 | 33 | 9-26 |
| HBK RY4721 | 66.5 | | | 1.6 | 1.6 | 36 | 9-17 |
| Asgrow AG 4632 | 65.5 | | | 1.6 | 1.3 | 32 | 9-18 |
| Asgrow AG 4832 | 64.9 | | | 2.3 | 1.9 | 36 | 9-19 |
| USG 74G99L | 64.3 | | | 1.4 | 1.1 | 35 | 9-25 |
| GoSoy 4910LL | 62.0 | | | 1.4 | 1.3 | 33 | 9-30 |
| HBK R4924 | 61.7 | | | 1.7 | 1.1 | 38 | 9-19 |
| USG 74H92R | 61.0 | | | 1.6 | 1.0 | 30 | 9-21 |
| Dyna-Gro S47RY13 | 59.7 | | | 1.3 | 1.9 | 30 | 9-17 |
| TN09-029 | 59.3 | | | 1.1 | 1.0 | 27 | 9-25 |
| GoSoy 4411LL | 58.9 | | | 1.9 | 3.0 | 32 | 9-15 |
| Asgrow AG 4633 | 54.1 | | | 1.3 | 1.6 | 26 | 9-12 |
| Terral REV 47R74 | 54.1 | | | 1.1 | 2.6 | 29 | 9-14 |
| Asgrow AG 4933 | 54.0 | | | 1.1 | 1.3 | 30 | 9-21 |
| Terral REV 49R54 | | | | 2.0 | 1.7 | 32 | 9-19 |
| Trial mean | 61.9 | 60.4 | 46.9 | 1.6 | 1.6 | 32 | 9-17 |
| LSD(0.10) | 4.8 | 4.4 | 46.9 | | | | |
| CV (%) | 10.8 | 13.8 | 45.5 | | | | |

[†] Multiyear averages do not include the trial at Belle Mina in 2011, which could not be planted due to excessive rain.

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 3. PERFORMANCE OF GROUP IV SOYBEAN VARIETIES IN TALLASSEE, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height | Maturity date |
|-------------------|----------------------|------------------|---------------------|-----------------|------------------|
| Maturity Group IV | | | | | |
| Asgrow AG 4933 | 63.4 | 1.0 | 1.8 | 34 | 9-16 |
| Asgrow AG 4832 | 63.0 | 1.5 | 2.0 | 38 | 9-14 |
| HBK R4924 | 60.4 | 1.3 | 1.8 | 43 | 9-16 |
| Terral REV 47R53 | 60.4 | 1.0 | 2.0 | 36 | 9-11 |
| Asgrow AG 4633 | 59.8 | 1.3 | 1.5 | 33 | 9-6 |
| Terral REV 49R22 | 56.8 | 2.3 | 2.5 | 38 | 9-16 |
| HBK RY4620 | 53.8 | 1.3 | 2.0 | 34 | 9-17 |
| Terral REV 49R43 | 53.1 | 1.0 | 2.3 | 35 | 9-12 |
| HBK RY4721 | 53.1 | 1.3 | 2.0 | 38 | 9-16 |
| Terral REV 47R74 | 52.4 | 1.0 | 2.3 | 36 | 9-5 |
| TN09-029 | 51.0 | 1.0 | 1.3 | 31 | 9-21 |
| Terral REV 48R33 | 49.3 | 1.0 | 2.0 | 38 | 9-14 |
| Trial mean | 56.4 | 1.2 | 1.9 | 36 | 9-14 |
| LSD(0.10) | 7.1 | | | | |
| CV (%) | 11.8 | | | | |

TABLE 4. PERFORMANCE OF GROUP IV SOYBEAN VARIETIES IN TALLASSEE, ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | | |
|-------------------|----------------|----------|----------|---------------|------------------|--------------|---------------|--|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | | |
| | <i>bu/acre</i> | | | | | | | | | |
| Maturity Group IV | | | | | | | | | | |
| Terral REV 49R22 | 56.8 | 54.4 | 49.7 | 1.4 | 2.0 | 34 | 9-8 | | | |
| Terral REV 47R53 | 60.4 | 50.7 | | 1.0 | 1.9 | 31 | 9-7 | | | |
| Terral REV 49R43 | 53.1 | 45.4 | | 1.0 | 1.9 | 29 | 9-6 | | | |
| Terral REV 48R33 | 49.3 | 43.2 | | 1.0 | 1.6 | 33 | 9-6 | | | |
| Asgrow AG 4933 | 63.4 | | | 1.0 | 1.8 | 34 | 9-16 | | | |
| Asgrow AG 4832 | 63.0 | | | 1.5 | 2.0 | 38 | 9-14 | | | |
| HBK R4924 | 60.4 | | | 1.3 | 1.8 | 43 | 9-16 | | | |
| Asgrow AG 4633 | 59.8 | | | 1.3 | 1.5 | 33 | 9-6 | | | |
| HBK RY4620 | 53.8 | | | 1.3 | 2.0 | 34 | 9-17 | | | |
| HBK RY4721 | 53.1 | | | 1.3 | 2.0 | 38 | 9-16 | | | |
| Terral REV 47R74 | 52.4 | | | 1.0 | 2.3 | 36 | 9-5 | | | |
| TN09-029 | 51.0 | | | 1.0 | 1.3 | 31 | 9-21 | | | |
| Trial mean | 56.4 | 48.4 | 49.7 | 1.2 | 1.8 | 34.3 | 9-11 | | | |
| LSD(0.10) | 7.1 | 3.7 | | | | | | | | |
| CV (%) | 11.8 | 13.4 | 1.7 | | | | | | | |

TABLE 5. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES IN NORTHERN ALABAMA, 2012

| Variety | Belle | Cross | Regional Average | | | | Maturity date |
|---------------------|-------|-------|------------------|------------------|---------------------|-------------------------------|------------------|
| | Mina | ville | Yield bu/acre | Lodging score | Shattering score | Plant height - inches - | |
| | ----- | ----- | | ----- | ----- | ----- | |
| Maturity Group IV | | | | | | | |
| Progeny P 4900RY | 68.7 | 57.0 | 62.8 | 1.2 | 1.1 | 35 | 10-2 |
| HBK RY4721 | 63.4 | 56.7 | 60.0 | 1.3 | 1.3 | 40 | 10-1 |
| Progeny P 4819LL | 52.8 | 54.2 | 53.5 | 1.6 | 1.1 | 37 | 10-3 |
| HBK R4924 | 54.4 | 49.7 | 52.1 | 2.0 | 1.4 | 43 | 10-2 |
| Progeny P 4920RY | 51.9 | 47.7 | 49.8 | 1.8 | 1.0 | 37 | 10-2 |
| Progeny P 4928LL | 55.3 | 43.8 | 49.5 | 1.3 | 1.3 | 37 | 10-1 |
| HBK RY4620 | 47.9 | 49.1 | 48.5 | 1.2 | 1.0 | 36 | 10-2 |
| Maturity Group V | | | | | | | |
| Progeny P 5210RY | 61.8 | 56.0 | 58.9 | 1.6 | 1.0 | 34 | 10-10 |
| Dyna-Gro 37RY52 | 55.3 | 57.6 | 56.5 | 1.4 | 1.0 | 34 | 10-10 |
| Dyna-Gro S54RY43 | 59.9 | 50.8 | 55.3 | 1.8 | 1.0 | 35 | 10-14 |
| Terral REV 53R23 | 57.2 | 52.7 | 55.0 | 1.1 | 1.1 | 29 | 10-7 |
| TN09-008 | 56.4 | 53.4 | 54.9 | 1.7 | 1.4 | 31 | 10-12 |
| Terral REV 54R84 | 56.3 | 53.3 | 54.8 | 2.3 | 1.0 | 30 | 10-8 |
| SS 5112NR2 | 60.4 | 49.3 | 54.8 | 1.3 | 1.0 | 33 | 10-11 |
| Terral REV 52R74 | 58.4 | 50.6 | 54.5 | 1.8 | 1.3 | 38 | 10-8 |
| Syngenta NK S 51-H9 | 59.1 | 49.2 | 54.2 | 1.3 | 1.0 | 35 | 10-10 |
| Progeny P 5711RY | 61.0 | 44.2 | 52.6 | 1.5 | 1.0 | 36 | 10-13 |
| Schillinger 5220.RC | 52.9 | 52.1 | 52.5 | 1.5 | 1.0 | 40 | 10-12 |
| Bayer HBK RY5221 | 56.9 | 47.7 | 52.3 | 2.0 | 1.0 | 40 | 10-9 |
| Progeny P 5412RY | 53.0 | 51.1 | 52.1 | 1.8 | 1.1 | 34 | 10-13 |
| Bayer HBK RY5421 | 56.4 | 45.4 | 50.9 | 1.7 | 1.1 | 31 | 10-8 |
| Terral REV 51R53 | 45.3 | 54.8 | 50.2 | 1.7 | 1.1 | 37 | 10-8 |
| Bayer HBK RY5425 | 50.5 | 47.5 | 49.0 | 2.6 | 1.0 | 43 | 10-15 |
| Progeny P 5460LL | 58.8 | 38.3 | 48.5 | 1.5 | 1.0 | 34 | 10-9 |
| Ozark | 56.1 | 40.8 | 48.5 | 1.6 | 1.4 | 34 | 10-10 |
| GoSoy 5410 LL | 50.2 | 42.0 | 46.1 | 1.0 | 1.1 | 37 | 10-3 |
| Progeny P 5111RY | 40.1 | 46.5 | 43.3 | 1.6 | 1.1 | 42 | 10-6 |
| Progeny P 5388RY | 55.9 | | | | | | |
| Trial mean | 55.6 | 49.7 | 52.6 | 1.6 | 1.1 | 36.1 | 10-8 |
| LSD(0.10) | 4.6 | 5.5 | 3.8 | | | | |
| CV (%) | 8.1 | 11.0 | 9.7 | | | | |

TABLE 6. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES IN NORTHERN ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | |
|---------------------|---------------------|-------------|-------------|---------------|------------------|--------------|---------------|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | |
| | ----- bu/acre ----- | | | | | - inch - | | | |
| Maturity Group IV | | | | | | | | | |
| Progeny P 4928LL | 49.5 | 45.6 | | 1.9 | 1.5 | 36 | 9-26 | | |
| Progeny P 4900RY | 62.8 | | | 1.1 | 1.1 | 35 | 10-2 | | |
| HBK RY4721 | 60.0 | | | 1.3 | 1.3 | 40 | 10-1 | | |
| Progeny P 4819LL | 53.5 | | | 1.6 | 1.1 | 36 | 10-3 | | |
| HBK R4924 | 52.1 | | | 2.0 | 1.4 | 43 | 10-2 | | |
| Progeny P 4920RY | 49.8 | | | 1.5 | 1.6 | 34 | 9-24 | | |
| HBK RY4620 | 48.5 | | | 1.1 | 1.0 | 35 | 10-3 | | |
| Maturity Group V | | | | | | | | | |
| Progeny P 5210RY | 58.9 | 52.7 | 41.7 | 2.4 | 1.4 | 33 | 9-29 | | |
| Ozark | 48.5 | 49.9 | 41.5 | 2.1 | 1.3 | 32 | 9-30 | | |
| Dyna-Gro 37RY52 | 56.5 | 52.2 | | 2.8 | 1.1 | 31 | 10-4 | | |
| Terral REV 51R53 | 50.2 | 51.5 | | 2.5 | 1.2 | 35 | 10-4 | | |
| SS 5112NR2 | 54.8 | 50.8 | | 2.1 | 1.3 | 33 | 10-6 | | |
| Schillinger 5220.RC | 52.5 | 47.5 | | 2.4 | 1.1 | 38 | 10-4 | | |
| Progeny P 5111RY | 43.3 | 45.9 | | 2.3 | 1.2 | 37 | 10-3 | | |
| Progeny P 5460LL | 48.5 | 45.5 | | 1.9 | 1.3 | 33 | 10-2 | | |
| Dyna-Gro S54RY43 | 55.3 | | | 1.9 | 1.0 | 35 | 10-14 | | |
| Terral REV 53R23 | 55.0 | | | 1.1 | 1.1 | 29 | 10-8 | | |
| TN09-008 | 54.9 | | | 1.7 | 1.4 | 31 | 10-12 | | |
| Terral REV 54R84 | 54.8 | | | 2.3 | 1.0 | 29 | 10-7 | | |
| Terral REV 52R74 | 54.5 | | | 1.7 | 1.3 | 38 | 10-9 | | |
| Syngenta NK S 51-H9 | 54.2 | | | 1.3 | 1.0 | 34 | 10-11 | | |
| Progeny P 5711RY | 52.6 | | | 1.4 | 1.0 | 35 | 10-13 | | |
| Bayer HBK RY5221 | 52.3 | | | 2.0 | 1.0 | 39 | 10-9 | | |
| Progeny P 5412RY | 52.1 | | | 1.9 | 1.1 | 34 | 10-13 | | |
| Bayer HBK RY5421 | 50.9 | | | 1.7 | 1.1 | 31 | 10-8 | | |
| Bayer HBK RY5425 | 49.0 | | | 2.7 | 1.0 | 43 | 10-15 | | |
| GoSoy 5410 LL | 46.1 | | | 1.0 | 1.1 | 37 | 10-4 | | |
| Progeny P 5388RY | | | | 2.7 | 1.0 | 32 | 10-11 | | |
| Trial mean | 52.6 | 49.1 | 41.6 | 1.9 | 1.2 | 35.0 | 10-6 | | |
| LSD(0.10) | 3.8 | 3.9 | 3.7 | | | | | | |
| CV (%) | 9.7 | 14.9 | 18.5 | | | | | | |

TABLE 7. PERFORMANCE OF MID-LATE GROUP V SOYBEAN VARIETIES IN NORTHERN ALABAMA, 2012

| Variety | Belle Mina | Cross ville | Yield | Regional Average | | | | |
|---------------------|---------------------|----------------|-------------|------------------|---------------------|-----------------|------------------|--|
| | | | | Lodging score | Shattering score | Plant height | Maturity date | |
| | ----- bu/acre ----- | | | | - inches - | | | |
| Maturity Group V | | | | | | | | |
| Terral REV 57R21 | 59.2 | 63.9 | 61.6 | 1.8 | 1.1 | 41 | 10-14 | |
| Progeny P 5160LL | 65.6 | 56.4 | 61.0 | 1.8 | 1.0 | 28 | 10-18 | |
| UA 5612 | 63.2 | 58.3 | 60.7 | 2.5 | 1.0 | 32 | 10-15 | |
| Asgrow AG 5831 | 65.0 | 54.8 | 59.9 | 1.0 | 1.0 | 29 | 10-13 | |
| USG Allen RR | 61.5 | 58.2 | 59.8 | 2.0 | 1.0 | 34 | 10-19 | |
| Progeny P 5210RY | 63.7 | 56.0 | 59.8 | 1.6 | 1.1 | 32 | 10-14 | |
| AGS 597RR | 62.8 | 56.4 | 59.6 | 2.0 | 1.0 | 33 | 10-20 | |
| Progeny P 5711RY | 61.3 | 57.9 | 59.6 | 2.1 | 1.1 | 33 | 10-13 | |
| SS 5911NR2 | 60.5 | 58.0 | 59.3 | 1.9 | 1.1 | 34 | 10-15 | |
| Asgrow AG 5533 | 63.4 | 54.2 | 58.8 | 2.8 | 1.1 | 32 | 10-13 | |
| Progeny P 5811RY | 59.1 | 57.9 | 58.5 | 2.0 | 1.1 | 34 | 10-17 | |
| Terral REV 55R83 | 61.6 | 55.0 | 58.3 | 1.8 | 1.1 | 34 | 10-16 | |
| Bayer HBK RY5521 | 58.8 | 57.7 | 58.2 | 2.1 | 1.0 | 33 | 10-17 | |
| Syngenta NK S 59-B8 | 61.4 | 54.9 | 58.2 | 2.5 | 1.1 | 37 | 10-15 | |
| Progeny P 5960LL | 62.4 | 53.2 | 57.8 | 2.0 | 1.1 | 34 | 10-17 | |
| Syngenta NK S 56-G6 | 60.6 | 53.9 | 57.3 | 1.1 | 1.3 | 32 | 10-13 | |
| Progeny P 5610RY | 58.5 | 56.1 | 57.3 | 2.4 | 1.0 | 33 | 10-17 | |
| Syngenta NK S 56-W5 | 61.7 | 52.8 | 57.2 | 2.4 | 1.0 | 35 | 10-19 | |
| Dyna-Gro 39RY57 | 56.6 | 57.4 | 57.0 | 1.8 | 1.0 | 30 | 10-14 | |
| Asgrow AG 5633 | 53.4 | 60.3 | 56.8 | 1.6 | 1.0 | 31 | 10-13 | |
| Osage | 57.7 | 55.9 | 56.8 | 1.1 | 1.0 | 29 | 10-12 | |
| Syngenta NK S 57-K3 | 59.8 | 53.7 | 56.7 | 2.1 | 1.0 | 35 | 10-15 | |
| Dyna-Gro 32RY55 | 57.2 | 55.7 | 56.5 | 1.4 | 1.0 | 33 | 10-15 | |
| Terral REV 56R63 | 57.8 | 54.3 | 56.0 | 2.4 | 1.0 | 37 | 10-18 | |
| Terral REV 59R13 | 54.0 | 57.4 | 55.7 | 2.1 | 1.0 | 38 | 10-17 | |
| Progeny P 5655RY | 56.9 | 53.2 | 55.0 | 2.3 | 1.0 | 39 | 10-15 | |
| Asgrow AG 5732 | 55.3 | 54.3 | 54.8 | 1.9 | 1.1 | 34 | 10-13 | |
| AGS 5911LL | 54.9 | 53.7 | 54.3 | 1.9 | 1.1 | 36 | 10-15 | |
| Terral REV 55R53 | 50.8 | 57.2 | 54.0 | 1.7 | 1.3 | 31 | 10-17 | |
| Terral REV 56R21 | 54.2 | 53.4 | 53.8 | 2.1 | 1.1 | 35 | 10-14 | |
| USG 75J62R | 56.6 | 50.9 | 53.7 | 2.9 | 1.1 | 41 | 10-16 | |
| SS 5510NR2 | 46.2 | 52.2 | 49.2 | 1.8 | 1.1 | 39 | 10-17 | |
| Trial mean | 58.8 | 55.8 | 57.3 | 2.0 | 1.1 | 33.9 | 10-15 | |
| LSD(0.10) | 4.1 | 4.9 | 3.2 | | | | | |
| CV (%) | 6.8 | 8.8 | 8.0 | | | | | |

**TABLE 8. PERFORMANCE OF MID-LATE GROUP V SOYBEAN VARIETIES IN NORTHERN ALABAMA,
THREE-YEAR SUMMARY, 2010 - 2012**

| Variety | Yield | | Lodging score | Shattering score | Plant height | Maturity date | | | |
|---------------------|---------------------|-------------|------------------|---------------------|-----------------|------------------|--|--|--|
| | 2012 | 2-yr avg | | | | | | | |
| | ----- bu/acre ----- | | | | | | | | |
| Maturity Group V | | | | | | | | | |
| Dyna-Gro 39RY57 | 57.0 | 58.3 | 2.9 | 1.1 | 32 | 10-9 | | | |
| Progeny P 5711RY | 59.6 | 57.4 | 2.6 | 1.4 | 35 | 10-10 | | | |
| Syngenta NK S 57-K3 | 56.7 | 56.8 | 2.6 | 1.1 | 36 | 10-12 | | | |
| AGS 597RR | 59.6 | 56.3 | 2.6 | 1.3 | 34 | 10-16 | | | |
| USG Allen RR | 59.8 | 56.0 | 2.1 | 1.2 | 36 | 10-15 | | | |
| Progeny P 5610RY | 57.3 | 56.0 | 2.6 | 1.0 | 34 | 10-8 | | | |
| Terral REV 56R21 | 53.8 | 54.7 | 2.8 | 1.3 | 36 | 10-8 | | | |
| Syngenta NK S 56-G6 | 57.3 | 54.5 | 2.1 | 1.4 | 34 | 10-8 | | | |
| Terral REV 56R63 | 56.0 | 54.2 | 2.6 | 1.3 | 37 | 10-11 | | | |
| Dyna-Gro 32RY55 | 56.5 | 54.1 | 2.1 | 1.1 | 35 | 10-9 | | | |
| Progeny P 5960LL | 57.8 | 53.8 | 2.1 | 1.1 | 35 | 10-9 | | | |
| Osage | 56.8 | 53.5 | 2.1 | 1.0 | 32 | 10-4 | | | |
| Progeny P 5811RY | 58.5 | 53.3 | 2.3 | 1.1 | 33 | 10-8 | | | |
| Terral REV 57R21 | 61.6 | 53.2 | 2.4 | 1.4 | 39 | 10-7 | | | |
| Progeny P 5655RY | 55.0 | 53.1 | 2.4 | 1.3 | 39 | 10-8 | | | |
| Asgrow AG 5831 | 59.9 | 52.3 | 1.6 | 1.1 | 32 | 10-6 | | | |
| Asgrow AG 5732 | 54.8 | 51.8 | 2.2 | 1.2 | 35 | 10-7 | | | |
| SS 5510NR2 | 49.2 | 46.5 | 2.2 | 1.4 | 40 | 10-8 | | | |
| Progeny P 5160LL | 61.0 | | 1.9 | 1.0 | 28 | 10-18 | | | |
| UA 5612 | 60.7 | | 2.6 | 1.0 | 32 | 10-15 | | | |
| Progeny P 5210RY | 59.8 | | 1.6 | 1.1 | 31 | 10-15 | | | |
| SS 5911NR2 | 59.3 | | 1.9 | 1.1 | 33 | 10-15 | | | |
| Asgrow AG 5533 | 58.8 | | 2.9 | 1.1 | 32 | 10-13 | | | |
| Terral REV 55R83 | 58.3 | | 1.9 | 1.1 | 34 | 10-16 | | | |
| Bayer HBK RY5521 | 58.2 | | 2.1 | 1.0 | 33 | 10-17 | | | |
| Syngenta NK S 59-B8 | 58.2 | | 2.6 | 1.1 | 37 | 10-15 | | | |
| Syngenta NK S 56-W5 | 57.2 | | 2.4 | 1.0 | 35 | 10-19 | | | |
| Asgrow AG 5633 | 56.8 | | 1.6 | 1.0 | 30 | 10-13 | | | |
| Terral REV 59R13 | 55.7 | | 2.1 | 1.0 | 38 | 10-18 | | | |
| AGS 5911LL | 54.3 | | 1.9 | 1.1 | 35 | 10-15 | | | |
| Terral REV 55R53 | 54.0 | | 1.7 | 1.3 | 31 | 10-17 | | | |
| USG 75J62R | 53.7 | | 3.0 | 1.1 | 42 | 10-16 | | | |
| Trial mean | 57.3 | 54.2 | 2.3 | 1.2 | 34.5 | 10-12 | | | |
| LSD(0.10) | 3.2 | 3.3 | | | | | | | |
| CV (%) | 8.0 | 11.7 | | | | | | | |

[†] Three-year averages not available; this is only the second year of this test.

**TABLE 9 PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES IN NORTHERN ALABAMA,
2012**

| Variety | Belle | Cross | Yield | Regional Average | | | Maturity date | |
|--------------------|---------------------|-------|-------|------------------|---------------------|-----------------|------------------|--|
| | Mina | ville | | Lodging score | Shattering score | Plant height | | |
| | ----- bu/acre ----- | | | - inches - | | | | |
| Maturity Group VI | | | | | | | | |
| Progeny P 6710RY | 60.3 | 59.2 | 59.7 | 2.1 | 2.1 | 37 | 10-17 | |
| R02-3065 | 57.5 | 53.5 | 55.5 | 1.6 | 1.6 | 31 | 10-15 | |
| R03-1250 | 55.5 | 53.2 | 54.3 | 2.1 | 2.1 | 36 | 10-16 | |
| AGS 6011LL | 54.4 | 52.0 | 53.2 | 2.0 | 2.0 | 32 | 10-14 | |
| SS RT 6810NR2 | 53.8 | 50.5 | 52.2 | 1.8 | 1.8 | 37 | 10-17 | |
| Maturity Group VII | | | | | | | | |
| Progeny P 7310RY | 56.5 | 54.4 | 55.4 | 2.0 | 2.0 | 32 | 10-17 | |
| Bayer HBK R7028 | 54.3 | 52.0 | 53.2 | 2.1 | 2.1 | 40 | 10-23 | |
| Bayer HBK R7200 | 34.0 | 56.1 | 45.0 | 2.8 | 2.8 | 40 | 10-15 | |
| Trial mean | 53.3 | 53.8 | 53.6 | 2.1 | 2.1 | 35.6 | 10-17 | |
| LSD(0.10) | 4.5 | 4.5 | | | | | | |
| CV (%) | 8.1 | 8.0 | | | | | | |

**TABLE 10. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES IN NORTHERN ALABAMA,
THREE-YEAR SUMMARY, 2010 - 2012**

| Variety | Yield | | | | | | Maturity date |
|--------------------|---------------------|----------|----------|---------------|------------------|--------------|---------------|
| | 2012 | 2-yr avg | 3-yr avg | Lodging score | Shattering score | Plant height | |
| | ----- bu/acre ----- | | | - inch - | | | |
| Maturity Group VI | | | | | | | |
| Progeny P 6710RY | 59.7 | 51.3 | 40.9 | 1.5 | 1.3 | 33 | 10-22 |
| SS RT 6810NR2 | 52.2 | 47.2 | 38.3 | 1.5 | 1.2 | 34 | 10-23 |
| R02-3065 | 55.5 | | | 1.6 | 1.0 | 30 | 10-15 |
| R03-1250 | 54.3 | | | 2.1 | 1.0 | 35 | 10-17 |
| AGS 6011LL | 53.2 | | | 2.0 | 1.0 | 32 | 10-14 |
| Maturity Group VII | | | | | | | |
| Progeny P 7310RY | 55.4 | 54.4 | 43.3 | 1.8 | 1.2 | 31 | 10-24 |
| Bayer HBK R7028 | 53.2 | | | 2.1 | 1.0 | 40 | 10-24 |
| Bayer HBK R7200 | 45.0 | | | 2.7 | 1.0 | 41 | 10-16 |
| Trial mean | 53.3 | 51.0 | 40.8 | 1.9 | 1.1 | 34.4 | 10-19 |
| LSD(0.10) | 4.5 | 4.5 | 3.1 | | | | |
| CV (%) | 8.1 | 16.0 | 17.0 | | | | |

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 11. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT SHORTER, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height | Maturity date |
|---------------------|----------------------|------------------|---------------------|-----------------|------------------|
| Maturity Group IV | | | | | |
| Progeny P 4920RY | 63.2 | 0.0 | -0.0 | 37 | 9-25 |
| Progeny P 4819LL | 62.3 | 0.0 | 0.0 | 36 | 9-25 |
| Progeny P 4900RY | 60.9 | 0.0 | 0.5 | 34 | 9-27 |
| Progeny P 4928LL | 58.1 | 0.0 | 0.5 | 38 | 9-27 |
| Maturity Group V | | | | | |
| Progeny P 5711RY | 66.0 | 0.0 | 0.0 | 37 | 10-3 |
| Progeny P 5160LL | 65.6 | 0.0 | 0.0 | 32 | 9-30 |
| Syngenta NK S 51-H9 | 65.1 | 0.0 | -0.0 | 36 | 9-26 |
| Progeny P 5412RY | 63.8 | 0.0 | 0.0 | 38 | 9-29 |
| TN09-008 | 63.5 | 0.0 | 0.0 | 35 | 10-2 |
| Bayer HBK RY5421 | 63.2 | 0.5 | 0.0 | 33 | 9-28 |
| Progeny P 5210RY | 63.1 | 0.0 | -0.0 | 38 | 9-28 |
| Terral REV 51R53 | 63.0 | 0.0 | 0.0 | 35 | 9-28 |
| Terral REV 53R23 | 62.9 | 0.0 | -0.0 | 32 | 9-25 |
| Ozark | 62.6 | 0.0 | 0.0 | 37 | 9-27 |
| SS 5112NR2 | 61.7 | 0.0 | 0.0 | 38 | 9-26 |
| Terral REV 52R74 | 60.3 | 0.0 | 0.3 | 41 | 9-30 |
| Progeny P 5460LL | 58.1 | -0.0 | 0.8 | 38 | 9-29 |
| Terral REV 54R84 | 56.6 | 3.3 | 0.0 | 34 | 9-28 |
| Progeny P 5111RY | 56.2 | 0.0 | 0.8 | 41 | 9-25 |
| Bayer HBK RY5221 | 55.9 | 0.0 | 0.5 | 43 | 9-25 |
| Progeny P 5388RY | 55.4 | 3.3 | 0.0 | 34 | 9-29 |
| Bayer HBK RY5425 | 54.6 | 0.3 | 0.0 | 43 | 10-6 |
| Trial mean | 61.0 | 0.3 | 0.1 | 37 | 9-28 |
| LSD(0.10) | 5.4 | | | | |
| CV (%) | 8.4 | | | | |

TABLE 12. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT SHORTER, ALABAMA, THREE-YEAR SUMMARY, 2010-2012

| Variety | Yield | | Lodging score | Shattering score | Plant height | Maturity date | | | |
|---------------------|---------|----------|---------------|------------------|--------------|---------------|--|--|--|
| | 2012 | 2-yr avg | | | | | | | |
| | bu/acre | | | | | | | | |
| Maturity Group IV | | | | | | | | | |
| Progeny P 4928LL | 58.1 | 38.6 | 0.0 | 0.5 | 37.8 | 9-25 | | | |
| Progeny P 4920RY | 63.2 | | 0.0 | 0.0 | 34.3 | 9-10 | | | |
| Progeny P 4819LL | 62.3 | | 0.0 | 0.0 | 36.0 | 9-25 | | | |
| Progeny P 4900RY | 60.9 | | 0.0 | 0.5 | 33.5 | 9-27 | | | |
| Maturity Group V | | | | | | | | | |
| Progeny P 5210RY | 63.1 | 46.7 | 44.4 | 0.0 | 35.8 | 9-17 | | | |
| Ozark | 62.6 | 46.5 | 43.5 | 0.0 | 32.4 | 9-19 | | | |
| Progeny P 5160LL | 65.6 | 52.2 | | 0.0 | 31.8 | 10-2 | | | |
| Progeny P 5711RY | 66.0 | 50.9 | | 0.0 | 34.5 | 10-5 | | | |
| Terral REV 51R53 | 63.0 | 48.9 | | 0.0 | 37.3 | 9-30 | | | |
| Progeny P 5111RY | 56.2 | 41.3 | | 0.8 | 39.1 | 9-24 | | | |
| Progeny P 5460LL | 58.1 | 40.0 | | 0.8 | 36.3 | 9-26 | | | |
| Syngenta NK S 51-H9 | 65.1 | | | 0.0 | 35.5 | 9-26 | | | |
| Progeny P 5412RY | 63.8 | | | 0.0 | 38.0 | 9-29 | | | |
| TN09-008 | 63.5 | | | 0.0 | 34.8 | 10-2 | | | |
| Bayer HBK RY5421 | 63.2 | | 0.5 | 0.0 | 33.3 | 9-28 | | | |
| Terral REV 53R23 | 62.9 | | 0.0 | 0.0 | 32.3 | 9-25 | | | |
| SS 5112NR2 | 61.7 | | 0.0 | 0.0 | 38.0 | 9-26 | | | |
| Terral REV 52R74 | 60.3 | | 0.0 | 0.3 | 41.3 | 9-30 | | | |
| Terral REV 54R84 | 56.6 | | 3.3 | 0.0 | 33.5 | 9-28 | | | |
| Bayer HBK RY5221 | 55.9 | | 0.0 | 0.5 | 42.8 | 9-25 | | | |
| Progeny P 5388RY | 55.4 | | 3.3 | 0.0 | 33.8 | 9-29 | | | |
| Bayer HBK RY5425 | 54.6 | | 0.3 | 0.0 | 43.3 | 10-6 | | | |
| Trial mean | 61.0 | 45.6 | 43.9 | 0.3 | 36.1 | 9-27 | | | |
| LSD(0.10) | 5.4 | 2.2 | 2.0 | | | | | | |
| CV (%) | 8.4 | 9.1 | 8.2 | | | | | | |

TABLE 13. PERFORMANCE PERFORMANCE OF MID-LATE GROUP V SOYBEAN VARIETIES OF SHORTER, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height | Maturity date |
|---------------------|----------------------|------------------|---------------------|--------------|---------------|
| | | - inches - | | | |
| Maturity Group V | | | | | |
| UA 5612 | 63.9 | 0.3 | 0.0 | 39.0 | 9-30 |
| Terral REV 55R53 | 62.6 | 0.0 | 0.0 | 35.5 | 10-1 |
| Progeny P 5711RY | 62.0 | 0.0 | 0.0 | 35.8 | 10-2 |
| Terral REV 57R21 | 61.7 | 0.0 | 0.3 | 39.5 | 9-28 |
| Terral REV 56R63 | 61.0 | 0.0 | 0.0 | 41.5 | 10-2 |
| Osage | 60.9 | 0.0 | 0.0 | 34.0 | 9-30 |
| Asgrow AG 5831 | 60.1 | 0.0 | 0.0 | 33.5 | 10-1 |
| Syngenta NK S 56-G6 | 59.2 | 0.0 | 0.0 | 35.5 | 9-30 |
| Terral REV 56R21 | 57.6 | 0.0 | 0.0 | 37.3 | 9-30 |
| Asgrow AG 5732 | 57.6 | 0.3 | 0.0 | 37.3 | 9-28 |
| Progeny P 5610RY | 57.1 | 0.0 | 0.3 | 36.8 | 10-1 |
| Syngenta NK S 57-K3 | 57.0 | 0.0 | 0.0 | 35.8 | 10-3 |
| Terral REV 55R83 | 56.7 | 0.0 | 0.0 | 41.8 | 9-28 |
| SS 5510NR2 | 56.7 | 0.0 | 0.0 | 40.8 | 9-28 |
| Progeny P 5210RY | 56.5 | 0.0 | 0.0 | 37.8 | 9-28 |
| Progeny P 5811RY | 56.3 | 0.0 | 0.0 | 35.8 | 9-28 |
| Bayer HBK RY5521 | 55.9 | 0.0 | 0.5 | 36.8 | 9-28 |
| USG 75J62R | 55.5 | 2.0 | 0.0 | 46.7 | 9-28 |
| USG Allen RR | 55.4 | 0.0 | 0.3 | 39.0 | 10-5 |
| Progeny P 5655RY | 55.3 | 0.0 | 0.0 | 41.5 | 9-28 |
| SS 5911NR2 | 55.3 | 0.0 | 0.3 | 35.5 | 10-6 |
| Syngenta NK S 59-B8 | 55.2 | 0.0 | 0.3 | 40.0 | 10-3 |
| Syngenta NK S 56-W5 | 54.9 | 0.0 | 0.0 | 40.0 | 9-30 |
| Progeny P 5960LL | 54.7 | 0.0 | 0.3 | 38.5 | 10-3 |
| Terral REV 59R13 | 54.0 | 0.0 | 0.0 | 37.3 | 10-3 |
| AGS 597RR | 53.2 | 0.0 | 0.0 | 38.3 | 10-5 |
| AGS 5911LL | 53.0 | 0.0 | 0.3 | 36.0 | 10-3 |
| Trial mean | 57.4 | 0.1 | 0.1 | 38 | 10-1 |
| LSD(0.10) | 5.9 | | | | |
| CV (%) | 9.8 | | | | |

TABLE 14. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES AT SHORTER, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging | Shattering | Plant height - inches - | Maturity date |
|-------------------|----------------------|------------|------------|----------------------------|---------------|
| | | score | score | | |
| Maturity Group V | | | | | |
| AGS 6011LL | 56.9 | 0.0 | 0.3 | 33.0 | 9-29 |
| R03-1250 | 56.9 | 0.0 | -0.0 | 31.0 | 10-3 |
| R02-3065 | 53.1 | 0.0 | 0.3 | 30.3 | 10-1 |
| SS RT 6810NR2 | 50.7 | 0.0 | -0.0 | 35.5 | 10-16 |
| Asgrow AG 6132 | 47.5 | 0.0 | 0.0 | 37.3 | 10-13 |
| Progeny P 6710RY | 47.2 | 0.0 | 0.0 | 36.0 | 10-16 |
| Woodruff | 52.7 | 0.0 | 0.0 | 38.5 | 10-21 |
| Progeny P 7310RY | 51.3 | 0.0 | 0.0 | 34.0 | 10-17 |
| AGS 787 RR | 46.4 | 0.0 | 0.0 | 38.5 | 10-15 |
| Bayer HBK R7028 | 45.8 | 0.0 | 0.0 | 36.8 | 10-13 |
| Bayer HBK R7200 | 41.5 | 0.3 | 0.3 | 46.5 | 10-9 |
| Henderson | 54.0 | 0.0 | 0.0 | 37.0 | 10-21 |
| Trial mean | 50.3 | 0.0 | 0.1 | 36.2 | 10-12 |
| LSD(0.10) | 4.9 | | | | |
| CV (%) | 9.2 | | | | |

TABLE 15. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES AT SHORTER, ALABAMA, THREE-YEAR SUMMARY, 2010-2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | | | |
|-------------------|---------------------|-------------|-------------|---------------|------------------|--------------|---------------|--|--|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | | | |
| | ----- bu/acre ----- | | | | | | | | | | |
| | Maturity Group VI | | | | | | | | | | |
| Progeny P 6710RY | 47.2 | 48.4 | 48.4 | 0.0 | 0.0 | 33.9 | 10-20 | | | | |
| Asgrow AG 6132 | 47.5 | 45.6 | 45.6 | 0.0 | 0.0 | 36.8 | 10-18 | | | | |
| AGS 6011LL | 56.9 | | | 0.0 | 0.3 | 33.0 | 9-29 | | | | |
| R03-1250 | 56.9 | | | 0.0 | 0.0 | 31.0 | 10-3 | | | | |
| R02-3065 | 53.1 | | | 0.0 | 0.3 | 30.3 | 10-1 | | | | |
| SS RT 6810NR2 | 50.7 | | | 0.0 | 0.0 | 35.5 | 10-16 | | | | |
| | Maturity Group VII | | | | | | | | | | |
| Woodruff | 52.7 | 53.2 | 53.2 | 0.0 | 0.0 | 35.9 | 10-24 | | | | |
| Progeny P 7310RY | 51.3 | 51.5 | 51.5 | 0.0 | 0.0 | 32.3 | 10-22 | | | | |
| AGS 787 RR | 46.4 | | | 0.0 | 0.0 | 38.5 | 10-15 | | | | |
| Bayer HBK R7028 | 45.8 | | | 0.0 | 0.0 | 36.8 | 10-13 | | | | |
| Bayer HBK R7200 | 41.5 | | | 0.3 | 0.3 | 46.5 | 10-9 | | | | |
| | Maturity Group VIII | | | | | | | | | | |
| Henderson | 54.0 | 54.4 | 54.4 | 0.0 | 0.0 | 35.8 | 10-22 | | | | |
| Trial mean | 50.3 | 50.6 | 50.6 | 0.0 | 0.1 | 35.5 | 10-14 | | | | |
| LSD(0.10) | 4.9 | 1.9 | 1.5 | | | | | | | | |
| CV (%) | 9.2 | 6.7 | 6.7 | | | | | | | | |

TABLE 16. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES ON VAIDEN SOIL, MARION JUNCTION, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging | Shattering | Plant | Maturity |
|-------------------|----------------------|------------|------------|-----------|-------------|
| | | score | score | height | date |
| Maturity Group IV | | | | | |
| Progeny P 4900RY | 52.6 | 1.3 | 1.5 | 31 | 9-25 |
| Progeny P 4928LL | 52.3 | 1.3 | 1.5 | 35 | 9-29 |
| Progeny P 4819LL | 50.8 | 1.0 | 1.0 | 31 | 9-26 |
| Progeny P 4920RY | 45.9 | 2.3 | 1.0 | 36 | 9-25 |
| Maturity Group V | | | | | |
| Dyna-Gro 39RY57 | 56.8 | 1.0 | 1.0 | 33 | 10-6 |
| Progeny P 5655RY | 55.2 | 1.5 | 1.0 | 37 | 10-7 |
| Progeny P 5160LL | 54.8 | 1.0 | 1.0 | 24 | 10-2 |
| Progeny P 5610RY | 54.5 | 1.3 | 1.0 | 32 | 10-4 |
| Progeny P 5811RY | 54.3 | 1.0 | 1.0 | 33 | 10-4 |
| Terral REV 56R21 | 54.3 | 1.3 | 1.0 | 31 | 10-4 |
| Terral REV 55R83 | 54.1 | 1.5 | 1.0 | 32 | 10-5 |
| Progeny P 5960LL | 53.9 | 1.0 | 1.0 | 33 | 10-7 |
| Progeny P 5210RY | 53.9 | 1.0 | 1.0 | 33 | 9-30 |
| Bayer HBK RY5221 | 52.5 | 2.5 | 1.0 | 38 | 9-30 |
| Bayer HBK RY5425 | 52.5 | 1.5 | 1.0 | 43 | 10-10 |
| Terral REV 51R53 | 52.5 | 2.5 | 1.0 | 36 | 10-3 |
| Bayer HBK RY5421 | 52.4 | 1.3 | 1.0 | 28 | 9-29 |
| Progeny P 5412RY | 52.1 | 1.3 | 1.0 | 32 | 10-2 |
| Asgrow AG 5831 | 52.0 | 1.0 | 1.0 | 29 | 10-4 |
| Terral REV 56R63 | 51.8 | 1.5 | 1.0 | 33 | 10-9 |
| Bayer HBK RY5521 | 51.7 | 1.0 | 1.0 | 29 | 10-3 |
| Terral REV 55R53 | 51.6 | 2.0 | 1.0 | 28 | 10-2 |
| Progeny P 5711RY | 51.5 | 1.0 | 1.0 | 30 | 10-4 |
| Progeny P 5111RY | 51.4 | 1.3 | 1.5 | 38 | 10-1 |
| Terral REV 59R13 | 50.3 | 1.0 | 1.0 | 37 | 10-7 |
| Progeny P 5460LL | 50.3 | 1.0 | 1.5 | 36 | 9-29 |
| Terral REV 54R84 | 49.3 | 1.5 | 2.0 | 26 | 9-30 |
| TN09-008 | 47.3 | 1.0 | 1.5 | 26 | 10-4 |
| Trial mean | 52.2 | 1.3 | 1.1 | 32 | 10-2 |
| LSD(0.10) | 5.2 | | | | |
| CV (%) | 9.5 | | | | |

TABLE 17. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES ON VAIDEN SOIL, MARION JUNCTION, ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | | | | Maturity date |
|-------------------|---------------------|----------|----------|---------------|------------------|--------------|---------------|
| | 2012 | 2-yr avg | 3-yr avg | Lodging score | Shattering score | Plant height | |
| | ----- bu/acre ----- | | | - inch - | | | |
| Maturity Group IV | | | | | | | |
| Progeny P 4928LL | 52.3 | 41.2 | | 1.1 | 3.0 | 35.0 | 9-22 |
| Progeny P 4900RY | 52.6 | | | 1.3 | 1.5 | 31.3 | 9-25 |
| Progeny P 4819LL | 50.8 | | | 1.0 | 1.0 | 30.5 | 9-26 |
| Progeny P 4920RY | 45.9 | | | 2.1 | 2.0 | 31.5 | 9-24 |
| Maturity Group V | | | | | | | |
| Progeny P 5610RY | 54.5 | 48.5 | 42.8 | 1.3 | 1.0 | 30.6 | 9-24 |
| Terral REV 56R21 | 54.3 | 46.6 | 41.4 | 1.3 | 1.2 | 30.4 | 9-22 |
| Progeny P 5210RY | 53.9 | 46.7 | 39.8 | 1.0 | 1.2 | 28.5 | 9-20 |
| Asgrow AG 5831 | 52.0 | 44.4 | 38.0 | 1.0 | 1.0 | 25.2 | 9-23 |
| Progeny P 5160LL | 54.8 | 48.3 | | 1.0 | 1.3 | 23.7 | 9-25 |
| Progeny P 5655RY | 55.2 | 47.7 | | 1.4 | 1.6 | 38.7 | 9-28 |
| Progeny P 5811RY | 54.3 | 46.8 | | 1.0 | 1.3 | 33.1 | 9-27 |
| Progeny P 5960LL | 53.9 | 46.6 | | 1.1 | 1.0 | 32.7 | 9-29 |
| Terral REV 56R63 | 51.8 | 46.5 | | 1.6 | 1.0 | 34.4 | 9-30 |
| Progeny P 5711RY | 51.5 | 46.1 | | 1.0 | 1.0 | 31.1 | 9-27 |
| Terral REV 51R53 | 52.5 | 46.1 | | 2.6 | 1.3 | 36.6 | 9-26 |
| Progeny P 5111RY | 51.4 | 43.3 | | 1.1 | 2.1 | 33.6 | 9-23 |
| Progeny P 5460LL | 50.3 | 40.5 | | 1.0 | 2.5 | 35.2 | 9-25 |
| Dyna-Gro 39RY57 | 56.8 | | | 1.0 | 1.0 | 33.3 | 10-6 |
| Terral REV 55R83 | 54.1 | | | 1.5 | 1.0 | 31.8 | 10-5 |
| Bayer HBK RY5221 | 52.5 | | | 2.5 | 1.0 | 38.0 | 9-30 |
| Bayer HBK RY5425 | 52.5 | | | 1.5 | 1.0 | 42.8 | 10-10 |
| Bayer HBK RY5421 | 52.4 | | | 1.3 | 1.0 | 27.5 | 9-29 |
| Progeny P 5412RY | 52.1 | | | 1.3 | 1.0 | 32.0 | 10-2 |
| Bayer HBK RY5521 | 51.7 | | | 1.0 | 1.0 | 28.5 | 10-3 |
| Terral REV 55R53 | 51.6 | | | 2.0 | 1.0 | 28.0 | 10-2 |
| Terral REV 59R13 | 50.3 | | | 1.0 | 1.0 | 37.3 | 10-7 |
| Terral REV 54R84 | 49.3 | | | 1.5 | 2.0 | 26.3 | 9-30 |
| TN09-008 | 47.3 | | | 1.0 | 1.5 | 26.0 | 10-4 |
| Trial mean | 52.2 | 45.7 | 40.5 | 1.3 | 1.3 | 31.9 | 9-28 |
| LSD(0.10) | 5.2 | 2.3 | 1.9 | | | | |
| CV (%) | 9.5 | 9.3 | 9.9 | | | | |

TABLE 18. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES ON SUMTER SOIL, MARION JUNCTION, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height - inches - | Maturity date | Iron Chlorosis [†] |
|-------------------|----------------------|------------------|---------------------|-------------------------------|------------------|--------------------------------|
| Maturity Group IV | | | | | | |
| Progeny P 4900RY | 26.4 | 1.0 | 1.5 | 21 | 10-14 | 2.3 |
| Progeny P 4928LL | 25.4 | 1.0 | 1.5 | 19 | 10-14 | 1.9 |
| Progeny P 4920RY | 20.9 | 1.0 | 1.0 | 20 | 10-9 | 3.6 |
| Progeny P 4819LL | 16.9 | 1.0 | 2.0 | 15 | 10-17 | 5.8 |
| Maturity Group V | | | | | | |
| Dyna-Gro 39RY57 | 38.9 | 1.0 | 2.0 | 19 | 10-3 | 1.3 |
| Progeny P 5655RY | 37.1 | 1.0 | 1.5 | 22 | 10-3 | 2.1 |
| Asgrow AG 5831 | 29.1 | 1.0 | 1.0 | 17 | 10-4 | 1.3 |
| Progeny P 5960LL | 28.8 | 1.0 | 1.0 | 22 | 10-4 | 1.0 |
| Progeny P 5210RY | 27.6 | 1.0 | 1.0 | 16 | 10-3 | 1.6 |
| Terral REV 56R63 | 26.6 | 1.0 | 1.0 | 18 | 10-12 | 3.9 |
| Progeny P 5610RY | 25.7 | 1.0 | 1.5 | 21 | 10-11 | 1.6 |
| Terral REV 51R53 | 25.0 | 1.0 | 1.0 | 22 | 10-15 | 4.8 |
| Progeny P 5160LL | 23.6 | 1.0 | 1.0 | 14 | 10-3 | 1.9 |
| Terral REV 59R13 | 23.3 | 1.0 | 1.0 | 17 | 10-13 | 4.8 |
| Progeny P 5711RY | 22.2 | 1.0 | 1.5 | 14 | 10-6 | 4.6 |
| Terral REV 55R83 | 21.0 | 1.0 | 1.5 | 18 | 10-11 | 3.8 |
| Terral REV 56R21 | 19.9 | 1.0 | 1.0 | 15 | 10-15 | 4.6 |
| Bayer HBK RY5221 | 18.9 | 1.0 | 1.0 | 18 | 10-19 | 5.1 |
| Bayer HBK RY5421 | 17.7 | 1.0 | 1.5 | 13 | 10-14 | 4.6 |
| Bayer HBK RY5521 | 17.6 | 1.0 | 2.0 | 17 | 10-17 | 4.8 |
| Bayer HBK RY5425 | 17.2 | 1.0 | 1.0 | 19 | 10-21 | 6.8 |
| Progeny P 5460LL | 14.7 | 1.0 | 2.8 | 17 | 10-16 | 4.6 |
| Progeny P 5811RY | 13.7 | 1.0 | 1.5 | 13 | 10-18 | 6.0 |
| Progeny P 5412RY | 13.4 | 1.0 | 1.0 | 14 | 10-14 | 5.6 |
| Terral REV 54R84 | 7.7 | 1.0 | 2.0 | 11 | 10-16 | 6.1 |
| Terral REV 55R53 | 4.9 | 1.0 | 3.0 | 12 | 10-19 | 5.8 |
| TN09-008 | 2.1 | 1.0 | 3.0 | 9 | 10-15 | 8.0 |
| Progeny P 5111RY | 0.0 | | | | | 9.3 |
| Trial mean | 20.2 | 1.0 | 1.5 | 17 | 10-12 | 4.2 |
| LSD(0.10) | 7.2 | | | | | |
| CV (%) | 33.6 | | | | | |

[†] Iron chlorosis ratings made on July 24, 2012. 1 = no chlorosis; 10 = plants losing leaves due to necrotic spots on leaves.

TABLE 19. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES ON SUMTER SOIL, MARION JUNCTION, ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | Iron Chlorosis | | | | |
|-------------------|---------------------|----------|----------|---------------|------------------|--------------|---------------|----------------|--|--|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | | | | |
| | ----- bu/acre ----- | | | | | | | | | | | |
| Maturity Group IV | | | | | | | | | | | | |
| Progeny P 4928LL | 25.4 | 16.4 | | 1.0 | 1.3 | 16.4 | 10-11 | 4.6 | | | | |
| Progeny P 4900RY | 26.4 | | | 1.0 | 1.5 | 20.5 | 10-14 | 2.3 | | | | |
| Progeny P 4920RY | 20.9 | | | 1.0 | 1.0 | 19.5 | 10-9 | 5.4 | | | | |
| Progeny P 4819LL | 16.9 | | | 1.0 | 2.0 | 15.0 | 10-17 | 5.8 | | | | |
| Maturity Group V | | | | | | | | | | | | |
| Progeny P 5210RY | 27.6 | 19.8 | | 1.0 | 1.3 | 14.3 | 9-30 | 4.7 | | | | |
| Asgrow AG 5831 | 29.1 | 18.8 | | 1.0 | 1.3 | 15.1 | 9-28 | 3.9 | | | | |
| Progeny P 5610RY | 25.7 | 19.8 | | 1.0 | 2.1 | 18.4 | 10-4 | 4.2 | | | | |
| Terral REV 56R21 | 19.9 | 14.8 | | 1.0 | 1.0 | 14.7 | 10-12 | 6.6 | | | | |
| Progeny P 5655RY | 37.1 | 27.4 | | 1.0 | 3.0 | 21.1 | 9-30 | 2.3 | | | | |
| Progeny P 5960LL | 28.8 | 22.7 | | 1.0 | 1.6 | 21.0 | 10-1 | 1.0 | | | | |
| Terral REV 51R53 | 25.0 | 19.0 | | 1.0 | 1.3 | 20.3 | 10-13 | 5.9 | | | | |
| Terral REV 56R63 | 26.6 | 18.8 | | 1.0 | 1.3 | 16.7 | 10-10 | 4.7 | | | | |
| Progeny P 5711RY | 22.2 | 17.3 | | 1.0 | 1.9 | 14.3 | 9-30 | 4.6 | | | | |
| Progeny P 5160LL | 23.6 | 16.1 | | 1.0 | 1.6 | 13.0 | 9-29 | 2.9 | | | | |
| Progeny P 5811RY | 13.7 | 10.2 | | 1.0 | 1.7 | 13.8 | 10-14 | 6.3 | | | | |
| Progeny P 5460LL | 14.7 | 9.9 | | 1.0 | 2.4 | 17.1 | 10-13 | 5.3 | | | | |
| Progeny P 5111RY | 0.0 | 1.8 | | 1.0 | 3.7 | 11.0 | 9-18 | 6.7 | | | | |
| Dyna-Gro 39RY57 | 38.9 | | | 1.0 | 2.0 | 18.8 | 10-3 | 1.3 | | | | |
| Terral REV 59R13 | 23.3 | | | 1.0 | 1.0 | 16.5 | 10-13 | 4.8 | | | | |
| Terral REV 55R83 | 21.0 | | | 1.0 | 1.5 | 18.3 | 10-11 | 3.8 | | | | |
| Bayer HBK RY5221 | 18.9 | | | 1.0 | 1.0 | 17.8 | 10-19 | 5.1 | | | | |
| Bayer HBK RY5421 | 17.7 | | | 1.0 | 1.5 | 12.8 | 10-14 | 4.6 | | | | |
| Bayer HBK RY5521 | 17.6 | | | 1.0 | 2.0 | 16.5 | 10-17 | 4.8 | | | | |
| Bayer HBK RY5425 | 17.2 | | | 1.0 | 1.0 | 19.0 | 10-21 | 6.8 | | | | |
| Progeny P 5412RY | 13.4 | | | 1.0 | 1.0 | 13.7 | 10-14 | 5.6 | | | | |
| Terral REV 54R84 | 7.7 | | | 1.0 | 2.0 | 11.3 | 10-16 | 6.1 | | | | |
| Terral REV 55R53 | 4.9 | | | 1.0 | 3.0 | 12.0 | 10-19 | 5.8 | | | | |
| TN09-008 | 2.1 | | | 1.0 | 3.0 | 9.0 | 10-15 | 8.0 | | | | |
| Trial mean | 20.2 | 16.6 | | 1.0 | 1.7 | 16.0 | 10-9 | 4.8 | | | | |
| LSD(0.10) | 7.2 | 3.4 | | | | | | | | | | |
| CV (%) | 33.6 | 38.0 | | | | | | | | | | |

TABLE 20. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES ON VAIDEN SOIL, MARION JUNCTION, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging | Shattering | Plant | Maturity |
|---------------------|----------------------|------------|------------|-----------|--------------|
| | | score | score | height | date |
| Maturity Group VI | | | | | |
| Asgrow AG 6132 | 56.2 | 1.0 | 1.0 | 38 | 10-12 |
| NK Brand S67-R6 | 56.2 | 1.0 | 1.0 | 37 | 10-17 |
| NK Brand S68-D4 | 54.6 | 2.5 | 1.0 | 36 | 10-12 |
| Dyna-Gro 36RY68 | 54.5 | 1.3 | 1.0 | 39 | 10-17 |
| Progeny P 6710RY | 54.2 | 1.0 | 1.0 | 38 | 10-17 |
| Asgrow AG 6732 | 47.9 | 1.0 | 1.0 | 36 | 10-16 |
| Maturity Group VII | | | | | |
| Bayer HBK R7028 | 61.1 | 1.0 | 1.0 | 36 | 10-18 |
| Progeny P 7310RY | 59.7 | 1.5 | 1.0 | 35 | 10-16 |
| NK Brand S74-M3 | 59.4 | 1.0 | 1.0 | 37 | 10-17 |
| Dyna-Gro 34RY75 | 56.6 | 1.8 | 1.0 | 38 | 10-18 |
| Dyna-Gro V76N9RR | 56.6 | 1.8 | 1.0 | 36 | 10-19 |
| NK Brand S77-T7 | 55.5 | 1.8 | 1.0 | 39 | 10-16 |
| AGS 787 RR | 55.2 | 1.5 | 1.0 | 36 | 10-16 |
| Woodruff | 52.5 | 2.3 | 1.0 | 39 | 10-17 |
| NK Brand S78-G6 | 49.8 | 1.5 | 1.0 | 40 | 10-16 |
| Bayer HBK R7200 | 43.0 | 1.3 | 1.0 | 45 | 10-14 |
| Maturity Group VIII | | | | | |
| Henderson | 57.1 | 1.3 | 1.0 | 37 | 10-19 |
| Trial mean | 54.7 | 1.4 | 1.0 | 38 | 10-16 |
| LSD(0.10) | 3.6 | | | | |
| CV (%) | 6.2 | | | | |

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 21. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES ON VAIDEN SOIL, MARION JUNCTION, ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | | |
|---------------------|---------------------|----------|----------|---------------|------------------|--------------|---------------|--|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | | |
| | ----- bu/acre ----- | | | | | | | | | |
| Maturity Group VI | | | | | | | | | | |
| Dyna-Gro 36RY68 | 54.5 | 49.9 | 42.3 | 1.1 | 1.4 | 34.5 | 10-13 | | | |
| Progeny P 6710RY | 54.2 | 49.1 | 41.5 | 1.2 | 1.0 | 34.8 | 10-13 | | | |
| NK Brand S68-D4 | 54.6 | 49.1 | | 1.9 | 1.0 | 35.9 | 10-10 | | | |
| Asgrow AG 6132 | 56.2 | 47.8 | | 1.1 | 1.3 | 37.0 | 10-10 | | | |
| Asgrow AG 6732 | 47.9 | 43.6 | | 1.1 | 1.3 | 34.3 | 10-14 | | | |
| NK Brand S67-R6 | 56.2 | | | 1.0 | 1.0 | 36.5 | 10-17 | | | |
| Maturity Group VII | | | | | | | | | | |
| Progeny P 7310RY | 59.7 | 53.2 | 45.6 | 1.3 | 1.2 | 31.3 | 10-12 | | | |
| Woodruff | 52.5 | 47.6 | 40.2 | 2.1 | 1.2 | 35.3 | 10-15 | | | |
| Dyna-Gro V76N9RR | 56.6 | 48.0 | 39.3 | 1.5 | 2.1 | 33.8 | 10-17 | | | |
| NK Brand S78-G6 | 49.8 | 43.7 | | 1.9 | 1.0 | 38.7 | 10-17 | | | |
| Bayer HBK R7028 | 61.1 | | | 1.0 | 1.0 | 35.8 | 10-18 | | | |
| NK Brand S74-M3 | 59.4 | | | 1.0 | 1.0 | 37.0 | 10-17 | | | |
| Dyna-Gro 34RY75 | 56.6 | | | 1.8 | 1.0 | 38.3 | 10-18 | | | |
| NK Brand S77-T7 | 55.5 | | | 1.8 | 1.0 | 38.8 | 10-16 | | | |
| AGS 787 RR | 55.2 | | | 1.5 | 1.0 | 36.0 | 10-16 | | | |
| Bayer HBK R7200 | 43.0 | | | 1.3 | 1.0 | 44.8 | 10-14 | | | |
| Maturity Group VIII | | | | | | | | | | |
| Henderson | 57.1 | 52.3 | 39.3 | 1.2 | 1.7 | 31.8 | 10-16 | | | |
| Trial mean | 54.7 | 48.4 | 41.4 | 1.4 | 1.2 | 36.1 | 10-15 | | | |
| LSD(0.10) | 3.6 | 2.0 | 2.5 | | | | | | | |
| CV (%) | 6.0 | 7.7 | 13.6 | | | | | | | |

PERFORMANCE OF SOYBEAN VARIETIES IN ALABAMA, 2012

TABLE 22. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES ON SUMTER SOIL, MARION JUNCTION, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height - inches - | Maturity date | Iron Chlorosis [†] |
|---------------------|----------------------|------------------|---------------------|-------------------------------|------------------|--------------------------------|
| Maturity Group VI | | | | | | |
| NK Brand S68-D4 | 44.3 | 1.0 | 1.0 | 23 | 10-14 | 1.1 |
| NK Brand S67-R6 | 42.1 | 1.0 | 1.0 | 23 | 10-18 | 2.5 |
| Asgrow AG 6732 | 39.5 | 1.0 | 1.0 | 21 | 10-17 | 1.1 |
| Progeny P 6710RY | 38.6 | 1.0 | 1.0 | 20 | 10-16 | 1.5 |
| Dyna-Gro 36RY68 | 35.4 | 1.0 | 1.0 | 19 | 10-16 | 1.3 |
| Asgrow AG 6132 | 23.6 | 1.0 | 2.0 | 16 | 10-17 | 3.5 |
| Maturity Group VII | | | | | | |
| NK Brand S78-G6 | 43.3 | 1.0 | 1.0 | 25 | 10-18 | 1.3 |
| Woodruff | 38.9 | 1.0 | 1.0 | 19 | 10-21 | 2.9 |
| Progeny P 7310RY | 38.0 | 1.0 | 1.0 | 17 | 10-18 | 2.4 |
| Dyna-Gro V76N9RR | 37.4 | 1.0 | 1.0 | 18 | 10-19 | 3.6 |
| NK Brand S77-T7 | 34.8 | 1.0 | 1.0 | 18 | 10-18 | 4.3 |
| AGS 787 RR | 33.2 | 1.0 | 1.0 | 19 | 10-21 | 3.8 |
| Dyna-Gro 34RY75 | 31.9 | 1.0 | 1.0 | 16 | 10-18 | 4.5 |
| Bayer HBK R7028 | 31.2 | 1.0 | 1.0 | 18 | 10-19 | 4.1 |
| NK Brand S74-M3 | 29.2 | 1.0 | 1.0 | 18 | 10-19 | 2.8 |
| Bayer HBK R7200 | 0.0 | | | | | 8.8 |
| Maturity Group VIII | | | | | | |
| Henderson | 35.7 | 1.0 | 1.0 | 20 | 10-19 | 4.0 |
| Trial mean | 33.9 | 1.0 | 1.1 | 19.3 | 10-18 | 3.1 |
| LSD(0.10) | 8.9 | | | | | |
| CV (%) | 24.8 | | | | | |

[†] Iron chlorosis ratings made on July 24, 2012. 1 = no chlorosis; 10 = plants losing leaves due to necrotic spots on leaves.

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 23. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES ON SUMTER SOIL, MARION JUNCTION, ALABAMA, THREE-YEAR SUMMARY, 2010 - 2012

| Variety | Yield | | Lodging score | Shattering score | Plant height | Maturity date | Iron Chlorosis | | |
|---------------------|----------------|-------------|------------------|---------------------|-----------------|------------------|-------------------|--|--|
| | 2012 | 2-yr avg | | | | | | | |
| | <i>bu/acre</i> | | <i>- inch -</i> | | | | | | |
| Maturity Group VI | | | | | | | | | |
| Progeny P 6710RY | 38.6 | 27.1 | 1.0 | 1.0 | 16.1 | 10-17 | 3.1 | | |
| Dyna-Gro 36RY68 | 35.4 | 26.3 | 1.0 | 1.3 | 17.1 | 10-16 | 4.3 | | |
| NK Brand S68-D4 | 44.3 | 33.1 | 1.0 | 1.0 | 20.4 | 10-15 | 1.1 | | |
| Asgrow AG 6732 | 39.5 | 30.7 | 1.0 | 2.4 | 17.9 | 10-18 | 1.1 | | |
| Asgrow AG 6132 | 23.6 | 16.9 | 1.0 | 2.3 | 16.0 | 10-19 | 5.6 | | |
| NK Brand S67-R6 | 42.1 | | 1.0 | 1.0 | 23.0 | 10-18 | 2.5 | | |
| Maturity Group VII | | | | | | | | | |
| Woodruff | 38.9 | 30.5 | 1.0 | 1.3 | 17.1 | 10-23 | 5.9 | | |
| Dyna-Gro V76N9RR | 37.4 | 29.9 | 1.0 | 1.9 | 17.3 | 10-21 | 5.9 | | |
| Progeny P 7310RY | 38.0 | 30.0 | 1.0 | 1.0 | 15.9 | 10-19 | 5.2 | | |
| NK Brand S78-G6 | 43.3 | 34.3 | 1.0 | 1.0 | 22.3 | 10-19 | 1.3 | | |
| NK Brand S77-T7 | 34.8 | | 1.0 | 1.0 | 18.0 | 10-18 | 4.3 | | |
| AGS 787 RR | 33.2 | | 1.0 | 1.0 | 18.8 | 10-21 | 3.8 | | |
| Dyna-Gro 34RY75 | 31.9 | | 1.0 | 1.0 | 16.0 | 10-18 | 4.5 | | |
| Bayer HBK R7028 | 31.2 | | 1.0 | 1.0 | 18.3 | 10-19 | 4.1 | | |
| NK Brand S74-M3 | 29.2 | | 1.0 | 1.0 | 17.5 | 10-19 | 2.8 | | |
| Bayer HBK R7200 | 0.0 | | | | | | 8.8 | | |
| Maturity Group VIII | | | | | | | | | |
| Henderson | 35.7 | 28.6 | 1.0 | 1.6 | 19.6 | 10-22 | 6.4 | | |
| Trial mean | 33.9 | 28.7 | 1.0 | 1.3 | 18.2 | 10-19 | 4.1 | | |
| LSD(0.10) | 5.2 | 2.9 | | | | | | | |
| CV (%) | 20.4 | 18.8 | | | | | | | |

PERFORMANCE OF SOYBEAN VARIETIES IN ALABAMA, 2012

29

TABLE 24. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT FAIRHOPE, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging | Shattering | Plant | Maturity |
|-------------------|----------------------|---------|------------|--------|----------|
| | | score | score | height | date |
| Maturity Group IV | | | | | |
| Progeny P 4928LL | 51.7 | 3.8 | 3.5 | 41 | 9-29 |
| Progeny P 4900RY | 46.4 | 3.0 | 2.5 | 33 | 10-3 |
| Progeny P 4920RY | 44.0 | 4.0 | 1.0 | 36 | 9-30 |
| Progeny P 4819LL | 43.4 | 3.8 | 1.8 | 34 | 10-1 |
| Maturity Group V | | | | | |
| Progeny P 5711RY | 59.0 | 2.5 | 1.8 | 32 | 10-7 |
| Terral REV 54R84 | 55.8 | 2.8 | 2.4 | 25 | 10-5 |
| Terral REV 51R53 | 55.3 | 3.8 | 1.3 | 38 | 10-7 |
| Progeny P 5460LL | 54.7 | 3.1 | 2.9 | 38 | 10-3 |
| Progeny P 5412RY | 54.4 | 2.8 | 1.5 | 33 | 10-6 |
| AGS 597RR | 54.3 | 2.5 | 1.0 | 31 | 10-9 |
| TN09-008 | 54.1 | 2.5 | 1.5 | 29 | 10-7 |
| Terral REV 56R21 | 52.6 | 2.5 | 1.8 | 26 | 10-7 |
| Terral REV 55R83 | 52.5 | 2.0 | 1.5 | 33 | 10-6 |
| Terral REV 55R53 | 52.2 | 3.0 | 2.0 | 30 | 10-6 |
| Progeny P 5610RY | 51.6 | 3.0 | 2.0 | 29 | 10-8 |
| Progeny P 5655RY | 51.2 | 2.8 | 2.5 | 35 | 10-6 |
| Dyna-Gro 39RY57 | 50.6 | 2.5 | 2.5 | 29 | 10-5 |
| Progeny P 5210RY | 50.0 | 2.5 | 1.3 | 26 | 10-6 |
| Terral REV 56R63 | 49.5 | 3.0 | 1.3 | 33 | 10-8 |
| Progeny P 5811RY | 47.6 | 2.5 | 2.3 | 28 | 10-7 |
| Progeny P 5111RY | 47.5 | 3.0 | 2.8 | 47 | 10-7 |
| Progeny P 5160LL | 45.1 | 2.3 | 1.3 | 24 | 10-8 |
| Terral REV 59R13 | 43.1 | 3.0 | 1.0 | 36 | 10-8 |
| Progeny P 5960LL | 37.7 | 2.5 | 2.3 | 28 | 10-10 |
| AGS 5911LL | 35.2 | 3.0 | 2.3 | 33 | 10-8 |
| Trial mean | 49.6 | 2.9 | 1.9 | 32 | 10-6 |
| LSD(0.10) | 8.5 | | | | |
| CV (%) | 16.3 | | | | |

ALABAMA AGRICULTURAL EXPERIMENT STATION

TABLE 25. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT FAIRHOPE, ALABAMA, THREE-YEAR SUMMARY, 2010-2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | | |
|-------------------|---------------------|-------------|-------------|------------------|---------------------|-----------------|------------------|--|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | | |
| | ----- bu/acre ----- | | | | | | | | | |
| Maturity Group IV | | | | | | | | | | |
| Progeny P 4928LL | 51.7 | 34.3 | | 3.4 | 3.0 | 34.4 | 10-5 | | | |
| Progeny P 4900RY | | 46.4 | | 3.0 | 2.5 | 32.5 | 10-3 | | | |
| Progeny P 4920RY | | 44.0 | | 3.1 | 1.6 | 34.1 | 9-14 | | | |
| Progeny P 4819LL | | 43.4 | | 3.8 | 1.8 | 34.3 | 10-1 | | | |
| Maturity Group V | | | | | | | | | | |
| Terral REV 56R21 | 52.6 | 46.0 | 50.2 | 2.5 | 1.5 | 29.3 | 9-20 | | | |
| Progeny P 5610RY | 51.6 | 40.1 | 46.7 | 2.7 | 2.0 | 28.5 | 9-25 | | | |
| Progeny P 5210RY | 50.0 | 37.9 | 43.9 | 2.3 | 1.5 | 24.9 | 9-24 | | | |
| Terral REV 51R53 | 55.3 | 44.7 | | 3.6 | 1.5 | 36.0 | 10-9 | | | |
| Progeny P 5160LL | 45.1 | 43.2 | | 2.1 | 1.4 | 22.0 | 10-8 | | | |
| Progeny P 5111RY | 47.5 | 41.4 | | 2.5 | 2.3 | 38.0 | 10-4 | | | |
| Progeny P 5655RY | 51.2 | 41.4 | | 2.4 | 2.1 | 32.5 | 10-7 | | | |
| Progeny P 5811RY | 47.6 | 40.7 | | 2.4 | 1.9 | 25.8 | 10-7 | | | |
| Progeny P 5460LL | 54.7 | 40.6 | | 2.9 | 2.8 | 34.1 | 10-6 | | | |
| Progeny P 5711RY | 59.0 | 40.2 | | 2.3 | 1.6 | 25.9 | 10-8 | | | |
| Terral REV 56R63 | 49.5 | 39.0 | | 2.9 | 1.4 | 29.4 | 10-10 | | | |
| Progeny P 5960LL | 37.7 | 36.0 | | 2.4 | 1.6 | 28.5 | 10-11 | | | |
| Terral REV 54R84 | 55.8 | | | 2.8 | 2.4 | 24.9 | 10-5 | | | |
| Progeny P 5412RY | 54.4 | | | 2.8 | 1.5 | 33.0 | 10-6 | | | |
| AGS 597RR | 54.3 | | | 2.5 | 1.0 | 31.3 | 10-9 | | | |
| TN09-008 | 54.1 | | | 2.5 | 1.5 | 28.5 | 10-7 | | | |
| Terral REV 55R83 | 52.5 | | | 2.0 | 1.5 | 32.5 | 10-6 | | | |
| Terral REV 55R53 | 52.2 | | | 3.0 | 2.0 | 30.3 | 10-6 | | | |
| Dyna-Gro 39RY57 | 50.6 | | | 2.5 | 2.5 | 29.0 | 10-5 | | | |
| Terral REV 59R13 | 43.1 | | | 3.0 | 1.0 | 35.8 | 10-8 | | | |
| AGS 5911LL | 35.2 | | | 3.0 | 2.3 | 32.8 | 10-8 | | | |
| Trial mean | 49.6 | 40.4 | 46.9 | 2.7 | 1.8 | 30.7 | 10-4 | | | |
| LSD(0.10) | 5.3 | 5.3 | 2.7 | | | | | | | |
| CV (%) | 14.3 | 24.6 | 12.2 | | | | | | | |

TABLE 26. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES AT FAIRHOPE, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height - inches - | Maturity date |
|---------------------|----------------------|------------------|---------------------|-------------------------------|------------------|
| Maturity Group VI | | | | | |
| Asgrow AG 6931 | 55.9 | 3.0 | 1.0 | 35 | 10-19 |
| Dyna-Gro 36RY68 | 55.0 | 2.0 | 1.0 | 31 | 10-19 |
| Progeny P 6710RY | 54.0 | 2.3 | 1.0 | 35 | 10-19 |
| Asgrow AG 6732 | 53.1 | 2.5 | 1.0 | 34 | 10-19 |
| AGS 6011LL | 48.2 | 2.5 | 1.8 | 26 | 10-18 |
| Maturity Group VII | | | | | |
| Progeny P 7310RY | 61.2 | 2.0 | 1.0 | 32 | 10-20 |
| Dyna-Gro 34RY75 | 58.9 | 2.0 | 1.0 | 34 | 10-19 |
| Asgrow AG 7733 | 58.5 | 2.3 | 1.0 | 34 | 10-20 |
| Asgrow AG 7231 | 58.4 | 2.0 | 1.0 | 31 | 10-19 |
| Dyna-Gro V76N9RR | 58.1 | 2.8 | 1.8 | 30 | 10-22 |
| Asgrow AG 7532 | 54.4 | 3.0 | 1.0 | 34 | 10-19 |
| Asgrow AG 7333 | 54.0 | 2.3 | 1.0 | 34 | 10-19 |
| AGS 787 RR | 51.0 | 3.0 | 1.0 | 32 | 10-19 |
| Woodruff | 51.0 | 3.0 | 1.0 | 33 | 10-19 |
| Maturity Group VIII | | | | | |
| Henderson | 56.4 | 3.0 | 1.0 | 34 | 10-24 |
| Trial mean | 55.2 | 2.5 | 1.1 | 33 | 10-19 |
| LSD(0.10) | 5.0 | | | | |
| CV (%) | 8.5 | | | | |

**TABLE 27. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES AT FAIRHOPE, ALABAMA,
THREE-YEAR SUMMARY, 2010-2012**

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | |
|---------------------|----------------|-------------|-------------|------------------|---------------------|-----------------|------------------|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | |
| | <i>bu/acre</i> | | | | | <i>- inch -</i> | | | |
| Maturity Group VI | | | | | | | | | |
| Asgrow AG 6931 | 55.9 | 48.8 | 53.4 | 2.6 | 1.0 | 33.8 | 10-4 | | |
| Progeny P 6710RY | 54.0 | 39.9 | 47.4 | 2.0 | 1.0 | 29.3 | 10-5 | | |
| Dyna-Gro 36RY68 | 55.0 | 39.9 | 46.8 | 1.8 | 1.0 | 28.7 | 10-4 | | |
| Asgrow AG 6732 | 53.1 | 47.5 | | 2.3 | 1.4 | 29.4 | 10-22 | | |
| AGS 6011LL | 48.2 | | | 2.5 | 1.8 | 26.3 | 10-18 | | |
| Maturity Group VII | | | | | | | | | |
| Woodruff | 51.0 | 53.1 | 53.4 | 3.2 | 1.0 | 31.1 | 10-3 | | |
| Progeny P 7310RY | 61.2 | 42.0 | 50.8 | 2.0 | 1.1 | 27.8 | 10-4 | | |
| Dyna-Gro V76N9RR | 58.1 | 43.9 | 50.5 | 2.5 | 1.3 | 32.2 | 10-2 | | |
| Asgrow AG 7231 | 58.4 | 38.6 | 47.5 | 2.2 | 1.0 | 27.8 | 10-2 | | |
| Asgrow AG 7532 | 54.4 | 46.9 | | 2.5 | 1.0 | 30.5 | 10-22 | | |
| Dyna-Gro 34RY75 | 58.9 | | | 2.0 | 1.0 | 33.8 | 10-19 | | |
| Asgrow AG 7733 | 58.5 | | | 2.3 | 1.0 | 34.0 | 10-20 | | |
| Asgrow AG 7333 | 54.0 | | | 2.3 | 1.0 | 33.5 | 10-19 | | |
| AGS 787 RR | 51.0 | | | 3.0 | 1.0 | 30.3 | 10-19 | | |
| Maturity Group VIII | | | | | | | | | |
| Henderson | 56.4 | 42.9 | 46.5 | 2.8 | 1.1 | 29.9 | 10-6 | | |
| Trial mean | 55.2 | 44.3 | 49.5 | 2.4 | 1.1 | 30.5 | 10-11 | | |
| LSD(0.10) | 5.0 | 5.5 | 4.4 | | | | | | |
| CV (%) | 8.5 | 23.2 | 19.9 | | | | | | |

TABLE 28. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT BREWTON, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height | Maturity date |
|-------------------|----------------------|------------------|---------------------|-----------------|------------------|
| | Maturity Group IV | | | - inches - | |
| Progeny P 4920RY | 53.2 | 1.0 | 0.0 | 28 | 9-30 |
| Progeny P 4928LL | 52.5 | 1.0 | 0.0 | 28 | 10-4 |
| Progeny P 4900RY | 52.0 | 1.0 | 0.3 | 26 | 9-30 |
| Progeny P 4819LL | 51.9 | 1.0 | 0.3 | 25 | 9-27 |
| Maturity Group V | | | | | |
| Terral REV 54R84 | 64.9 | 1.0 | -0.0 | 24 | 10-12 |
| Terral REV 55R53 | 62.9 | 1.0 | 0.3 | 21 | 10-9 |
| Progeny P 5711RY | 61.6 | 1.0 | -0.0 | 24 | 10-13 |
| AGS 5911LL | 61.5 | 1.0 | -0.0 | 25 | 10-13 |
| Progeny P 5412RY | 61.0 | 1.0 | -0.0 | 27 | 10-10 |
| Progeny P 5210RY | 60.9 | 1.0 | -0.0 | 26 | 10-8 |
| UA 5612 | 60.5 | 1.0 | -0.0 | 26 | 10-10 |
| Progeny P 5160LL | 60.3 | 1.0 | -0.0 | 20 | 10-9 |
| Progeny P 5111RY | 59.0 | 1.0 | 0.3 | 34 | 10-6 |
| Progeny P 5960LL | 58.2 | 1.0 | -0.0 | 24 | 10-13 |
| Terral REV 56R63 | 57.3 | 1.0 | -0.0 | 27 | 10-13 |
| Progeny P 5811RY | 56.7 | 1.0 | -0.0 | 26 | 10-10 |
| Terral REV 51R53 | 56.7 | 1.0 | -0.0 | 31 | 10-6 |
| TN09-008 | 56.0 | 1.0 | 0.7 | 22 | 10-9 |
| Progeny P 5655RY | 54.3 | 1.0 | -0.0 | 29 | 10-10 |
| Terral REV 59R13 | 54.3 | 1.0 | -0.0 | 26 | 10-10 |
| Progeny P 5460LL | 54.3 | 1.0 | 0.2 | 29 | 10-7 |
| Progeny P 5610RY | 54.0 | 1.0 | 0.3 | 26 | 10-10 |
| Terral REV 55R83 | 53.3 | 1.0 | -0.0 | 25 | 10-12 |
| Terral REV 56R21 | 52.9 | 1.0 | -0.0 | 25 | 10-8 |
| AGS 597RR | 52.4 | 1.0 | -0.0 | 27 | 10-12 |
| Ozark | 52.0 | 1.0 | 0.0 | 23 | 10-7 |
| Osage | 51.1 | 1.0 | 0.0 | 22 | 10-10 |
| Trial mean | 56.5 | 1.0 | 0.1 | 26 | 10-8 |
| LSD(0.10) | 3.7 | | | | |
| CV (%) | 6.2 | | | | |

TABLE 29. PERFORMANCE OF GROUP IV AND V SOYBEAN VARIETIES AT BREWTON, ALABAMA, THREE-YEAR SUMMARY, 2010-2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | |
|-------------------|---------------------|-------------|-------------|------------------|---------------------|-----------------|------------------|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | |
| | ----- bu/acre ----- | | | | | - inch - | | | |
| Maturity Group IV | | | | | | | | | |
| Progeny P 4928LL | 52.5 | 34.9 | | 1.0 | 0.0 | 29.0 | 9-23 | | |
| Progeny P 4920RY | 53.2 | | | 1.7 | 3.3 | 30.7 | 9-13 | | |
| Progeny P 4900RY | 52.0 | | | 1.0 | 0.3 | 25.7 | 9-30 | | |
| Progeny P 4819LL | 51.9 | | | 1.0 | 0.3 | 25.3 | 9-27 | | |
| Maturity Group VI | | | | | | | | | |
| Progeny P 5210RY | 60.9 | 42.4 | 46.3 | 1.0 | 0.0 | 25.0 | 9-18 | | |
| Progeny P 5610RY | 54.0 | 43.0 | 45.8 | 1.0 | 0.2 | 24.9 | 9-18 | | |
| Terral REV 56R21 | 52.9 | 38.5 | 43.5 | 1.0 | 0.0 | 25.0 | 9-16 | | |
| Ozark | 52.0 | 38.8 | 43.0 | 1.0 | 0.0 | 23.0 | 9-16 | | |
| Osage | 51.1 | 34.5 | 35.8 | 1.0 | 1.7 | 21.8 | 9-16 | | |
| Terral REV 56R63 | 57.3 | 49.8 | | 1.0 | 0.0 | 25.3 | 10-3 | | |
| Progeny P 5711RY | 61.6 | 45.3 | | 1.0 | 0.0 | 23.8 | 10-2 | | |
| Progeny P 5160LL | 60.3 | 44.7 | | 1.0 | 0.0 | 19.8 | 9-27 | | |
| Progeny P 5111RY | 59.0 | 43.0 | | 1.0 | 0.3 | 28.0 | 9-25 | | |
| Progeny P 5960LL | 58.2 | 42.2 | | 1.0 | 0.0 | 25.3 | 10-2 | | |
| Terral REV 51R53 | 56.7 | 41.9 | | 1.0 | 0.0 | 34.7 | 9-26 | | |
| Progeny P 5811RY | 56.7 | 40.8 | | 1.0 | 0.0 | 26.0 | 9-28 | | |
| Progeny P 5655RY | 54.3 | 39.3 | | 1.0 | 0.0 | 26.8 | 9-28 | | |
| AGS 597RR | 52.4 | 37.6 | | 1.0 | 0.0 | 25.5 | 10-2 | | |
| Progeny P 5460LL | 54.3 | 37.5 | | 1.0 | 0.2 | 30.3 | 9-29 | | |
| Terral REV 54R84 | 64.9 | | | 1.0 | 0.0 | 24.3 | 10-12 | | |
| Terral REV 55R53 | 62.9 | | | 1.0 | 0.3 | 21.3 | 10-9 | | |
| AGS 5911LL | 61.5 | | | 1.0 | 0.0 | 25.3 | 10-13 | | |
| Progeny P 5412RY | 61.0 | | | 1.0 | 0.0 | 26.7 | 10-10 | | |
| UA 5612 | 60.5 | | | 1.0 | 0.0 | 25.7 | 10-10 | | |
| TN09-008 | 56.0 | | | 1.0 | 0.7 | 22.3 | 10-9 | | |
| Terral REV 59R13 | 54.3 | | | 1.0 | 0.0 | 26.3 | 10-10 | | |
| Terral REV 55R83 | 53.3 | | | 1.0 | 0.0 | 24.7 | 10-12 | | |
| Trial mean | 56.5 | 40.9 | 42.9 | 1.0 | 0.3 | 25.7 | 9-29 | | |
| LSD(0.10) | 2.4 | 3.5 | 2.3 | | | | | | |
| CV (%) | 5.7 | 15.9 | 11.4 | | | | | | |

TABLE 30. PERFORMANCE OF GROUP VI AND VII SOYBEAN VARIETIES AT BREWTON, ALABAMA, 2012

| Variety | Yield - bu/acre - | Lodging score | Shattering score | Plant height - inches - | Maturity date |
|---------------------|----------------------|------------------|---------------------|-------------------------------|------------------|
| Maturity Group VI | | | | | |
| Asgrow AG 6732 | 67.5 | 1.0 | 0.0 | 29 | 10-25 |
| Progeny P 6710RY | 64.3 | 1.0 | 0.0 | 24 | 10-18 |
| NK Brand S67-R6 | 63.2 | 1.0 | 0.0 | 30 | 10-19 |
| NK Brand S68-D4 | 58.7 | 1.0 | 0.0 | 29 | 10-14 |
| Asgrow AG 6931 | 56.8 | 1.0 | -0.0 | 31 | 10-20 |
| AGS 6011LL | 53.6 | 1.0 | 0.7 | 23 | 10-4 |
| R03-1250 | 51.5 | 1.0 | -0.0 | 21 | 10-9 |
| R02-3065 | 51.2 | 1.0 | 0.3 | 20 | 10-9 |
| Maturity Group VII | | | | | |
| NK Brand S77-T7 | 70.9 | 1.0 | 0.0 | 28 | 10-21 |
| NK Brand S74-M3 | 68.5 | 1.0 | 0.0 | 27 | 10-20 |
| Progeny P 7310RY | 65.4 | 1.0 | 0.0 | 24 | 10-19 |
| Asgrow AG 7733 | 63.4 | 1.0 | 0.0 | 31 | 10-24 |
| Asgrow AG 7231 | 63.0 | 1.0 | 0.0 | 26 | 10-20 |
| NK Brand S78-G6 | 61.9 | 1.0 | 0.0 | 32 | 10-22 |
| Asgrow AG 7333 | 61.0 | 1.0 | 0.0 | 29 | 10-20 |
| Asgrow AG 7532 | 60.3 | 1.0 | 0.0 | 27 | 10-19 |
| Maturity Group VIII | | | | | |
| Henderson | 59.5 | 1.0 | -0.0 | 31 | 10-24 |
| Trial mean | 61.2 | 1.0 | 0.1 | 27 | 10-18 |
| LSD(0.10) | 3.4 | | | | |
| CV (%) | 5.2 | | | | |

TABLE 31. PERFORMANCE OF SOYBEAN VARIETIES AT BREWTON, ALABAMA, THREE-YEAR SUMMARY, 2010-2012

| Variety | Yield | | | Lodging score | Shattering score | Plant height | Maturity date | | |
|---------------------|---------------------|-------------|-------------|------------------|---------------------|-----------------|------------------|--|--|
| | 2012 | 2-yr avg | 3-yr avg | | | | | | |
| | ----- bu/acre ----- | | | | | - inch - | | | |
| Maturity Group VI | | | | | | | | | |
| Asgrow AG 6931 | 56.8 | 49.5 | 48.6 | 1.0 | 0.0 | 31.8 | 9-30 | | |
| Progeny P 6710RY | 64.3 | 50.0 | 46.9 | 1.0 | 0.0 | 27.6 | 10-1 | | |
| Asgrow AG 6732 | 67.5 | 51.3 | | 1.0 | 0.0 | 27.3 | 10-21 | | |
| NK Brand S68-D4 | 58.7 | 46.7 | | 1.0 | 0.0 | 30.7 | 10-10 | | |
| NK Brand S67-R6 | 63.2 | | | 1.0 | 0.0 | 30.0 | 10-19 | | |
| AGS 6011LL | 53.6 | | | 1.0 | 0.7 | 23.3 | 10-4 | | |
| R03-1250 | 51.5 | | | 1.0 | 0.0 | 20.7 | 10-9 | | |
| R02-3065 | 51.2 | | | 1.0 | 0.3 | 20.0 | 10-9 | | |
| Maturity Group VII | | | | | | | | | |
| Progeny P 7310RY | 65.4 | 52.7 | 53.4 | 1.0 | 0.0 | 26.4 | 10-2 | | |
| Asgrow AG 7231 | 63.0 | 48.9 | 47.2 | 1.0 | 0.0 | 28.0 | 9-30 | | |
| NK Brand S78-G6 | 61.9 | 50.5 | | 1.0 | 0.0 | 35.3 | 10-19 | | |
| Asgrow AG 7532 | 60.3 | 48.3 | | 1.0 | 0.0 | 28.2 | 10-18 | | |
| NK Brand S77-T7 | 70.9 | | | 1.0 | 0.0 | 28.3 | 10-21 | | |
| NK Brand S74-M3 | 68.5 | | | 1.0 | 0.0 | 26.7 | 10-20 | | |
| Asgrow AG 7733 | 63.4 | | | 1.0 | 0.0 | 31.0 | 10-24 | | |
| Asgrow AG 7333 | 61.0 | | | 1.0 | 0.0 | 29.0 | 10-20 | | |
| Maturity Group VIII | | | | | | | | | |
| Henderson | 59.5 | 53.8 | 50.1 | 1.0 | 0.0 | 34.9 | 10-3 | | |
| Trial mean | 61.2 | 50.2 | 49.2 | 1.0 | 0.1 | 28.2 | 10-12 | | |
| LSD(0.10) | 3.4 | 2.7 | 2.6 | | | | | | |
| CV (%) | 5.2 | 9.9 | 11.4 | | | | | | |

TABLE 32. CULTURAL PRACTICES FOR SOYBEAN VARIETY TESTS IN 2012

| Location | Type of test | Date planted | Row width | Herbicide applied | Fertilizer applied |
|-----------------|-----------------------|--------------|-----------|--------------------|--------------------|
| - inches - | | | | | |
| Belle Mina | Group IV | April 25 | 30 | Reflex, Select Max | none recommended |
| | Group IV-V | June 7 | 30 | Valor | none recommended |
| | Group Mid-Late V | June 7 | 30 | Valor | none recommended |
| | Group VI-VII | June 7 | 30 | Valor | none recommended |
| Crossville | Group IV | April 25 | 30 | Dual, Valor | none recommended |
| | Group IV-V | May 29 | 30 | Dual, Valor | none recommended |
| | Group Mid-Late V | June 6 | 30 | Dual, Valor | none recommended |
| | Group VI-VII | June 7 | 30 | Dual, Valor | none recommended |
| Tallassee | Group IV | May 8 | 30 | None | none recommended |
| Shorter | Group IV-V | June 7 | 36 | Dual, Prowl | none recommended |
| | Group VI-VII | June 7 | 36 | Dual, Prowl | none recommended |
| Marion Junction | Group IV-V (Sumter) | June 4 | 36 | Basagram, Poast | none recommended |
| | Group VI-VII (Sumter) | June 4 | 36 | Basagram, Poast | none recommended |
| | Group IV-V (Vaiden) | June 4 | 36 | Basagram, Poast | none recommended |
| | Group VI-VII (Vaiden) | June 4 | 36 | Basagram, Poast | none recommended |
| Brewton | Group IV-V | June 20 | 36 | Dual | none recommended |
| | Group VI-VII | June 20 | 36 | Dual | none recommended |
| Fairhope | Group IV-V | May 31 | 38 | Dual, Storm | none recommended |
| | Group VI-VII | May 31 | 38 | Dual, Storm | none recommended |

TABLE 33. SOIL TYPES FOR SOYBEAN TESTS, 2012

| Location | Soil Type |
|-----------------|----------------------------|
| Belle Mina | Emory silt loam |
| Crossville | Wynnvile fine sandy loam |
| Tallassee | Augusta silt loam |
| Shorter | Cowarts loamy sand |
| Marion Junction | Sumter clay (high pH soil) |
| Marion Junction | Vaiden clay |
| Fairhope | Malbis fine sandy loam |

PERFORMANCE OF SOYBEAN VARIETIES IN ALABAMA, 2012

39

TABLE 34. RAINFALL AT TEST LOCATIONS DURING GROWING SEASON, 2012

| Month | Days | Belle Mina | Crossville | Shorter | Tallassee | Marion Junction | Brewton | Fairhope |
|---------------|-------|------------|------------|---------|-----------|-----------------|---------|----------|
| <i>inches</i> | | | | | | | | |
| May | 1-5 | 0.7 | 0.2 | 3.2 | 1.6 | 0.2 | 2.2 | 6.7 |
| | 6-10 | 0.1 | 0.4 | 1.0 | 0.7 | 0.7 | 0.2 | 0.3 |
| | 11-15 | 2.1 | 0.9 | 2.5 | 3.4 | 1.1 | 1.1 | 0.2 |
| | 16-20 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 21-25 | 0.2 | 0.5 | 1.6 | 0.2 | 1.6 | 0.4 | 0.0 |
| | 26-31 | 0.2 | 0.1 | 0.9 | 1.8 | 0.2 | 0.0 | 0.4 |
| June | 1-5 | 0.9 | 1.0 | 0.3 | 0.3 | 0.1 | 0.5 | 1.3 |
| | 6-10 | 0.1 | 0.7 | 0.8 | 0.7 | 0.2 | 3.2 | 6.7 |
| | 11-15 | 0.4 | 1.7 | 1.3 | 1.9 | 1.5 | 6.4 | 3.2 |
| | 16-20 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 2.0 | 0.0 |
| | 21-25 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 26-31 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| July | 1-5 | 0.4 | 0.7 | 0.9 | 0.5 | 0.4 | 0.7 | 1.0 |
| | 6-10 | 0.6 | 0.9 | 1.0 | 1.2 | 0.7 | 0.0 | 0.1 |
| | 11-15 | 5.1 | 1.8 | 1.7 | 1.0 | 4.7 | 7.8 | 1.9 |
| | 16-20 | 1.1 | 1.1 | 0.1 | 0.2 | 0.5 | 0.7 | 3.0 |
| | 21-25 | 0.4 | 1.3 | 0.2 | 0.0 | 0.1 | 0.0 | 1.1 |
| | 26-31 | 1.2 | 0.0 | 0.5 | 1.0 | 1.7 | 0.3 | 0.9 |
| August | 1-5 | 0.3 | 0.0 | 0.5 | 0.7 | 1.7 | 2.2 | 0.3 |
| | 6-10 | 0.1 | 2.8 | 2.5 | 1.2 | 0.8 | 1.2 | 1.2 |
| | 11-15 | 1.3 | 0.8 | 0.2 | 0.2 | 0.8 | 2.2 | 0.6 |
| | 16-20 | 1.8 | 0.6 | 1.5 | 1.7 | 2.4 | 3.5 | 2.4 |
| | 21-25 | 0.0 | 0.0 | 0.0 | 0.0 | 0.1 | 0.0 | 0.0 |
| | 26-31 | 0.0 | 0.0 | 0.2 | 0.1 | 0.8 | 2.4 | 0.0 |
| September | 1-5 | 1.2 | 3.7 | 1.4 | 1.1 | 5.4 | 1.9 | 1.5 |
| | 6-10 | 0.2 | 0.0 | 0.0 | 0.5 | 0.0 | 0.1 | 0.0 |
| | 11-15 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 16-20 | 4.1 | 1.8 | 0.4 | 0.5 | 1.6 | 2.0 | 1.2 |
| | 21-25 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 26-31 | 0.0 | 0.0 | 0.9 | 1.1 | 0.3 | 0.2 | 1.1 |
| October | 1-5 | 1.5 | 2.5 | 1.4 | 1.1 | 1.5 | 2.1 | 2.8 |
| | 6-10 | 0.4 | 0.1 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 11-15 | 1.2 | 0.3 | 0.9 | 0.9 | 0.2 | 0.0 | 0.0 |
| | 16-20 | 0.6 | 0.4 | 0.1 | 0.1 | 0.4 | 2.8 | 0.1 |
| | 21-25 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| | 26-31 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |

